For Research Use Only

C10orf67 Polyclonal antibody

Catalog Number:24459-1-AP



Basic Information	Catalog Number: 24459-1-AP	GenBank Accession Number: BC035732	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 750 ug/ml by Nanodrop and 347 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 256815 UNIPROT ID: O8IYJ2	Recommended Dilutions: WB 1:200-1:1000
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG21465	Full Name: chromosome 10 open reading frame 67 Calculated MW: 184 aa, 21 kDa Observed MW: 18-21 kDa	
Applications	Tested Applications: WB, ELISA Species Specificity: human	Positive Controls: WB : PC-3 cells, A549 cells, human testis tissue	
Background Information	RNF17 also termed as ring finger protein 17 is a 1623 amino acid protein, which contains 1 ring type zinc finger and 4 tudor domains. RNF17 is specifically expressed in testis. The RING finger motif presents in many ubiquitin E3 ligases. RNF17 interacts with all four members of the Mad family (Mad1, Mxi1, Mad3 and Mad4), which are basic-helix-loop-helix-leucine zipper transcription factors of the Myc oncoprotein network. RNF17 is component of the tectonic-like complex, which is required for ciliogenesis and sonic hedgehog/SHH signaling. RNF17 is able to form dimers or polymers both in vitro and in vivo, indicating that it may play a role in the assembly of RNF17 granules. RNF17 encodes a novel key regulator of spermiogenesis.		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3.		
*** 20ul sizes contain 0.1% BSA	Aliquoting is unnecessary for -20 $^{\circ}$ C s	torage	

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



PC-3 cells were subjected to SDS PAGE followed by western blot with 24459-1-AP (C10orf67 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.