For Research Use Only

SALL4 Polyclonal antibody

Catalog Number:24500-1-AP

Featured Product

8 Publications



Basic Information

Catalog Number: GenBank Accession Number: 24500-1-AP BC111714

Size: Genel D (NCBI):

150ul , Concentration: 450 μ g/ml by 57167

Nanodrop and 373 µg/ml by Bradford Full Name:

method using BSA as the standard; sal-like 4 (Drosophila)

Source: Calculated MW:
Rabbit 1053 aa, 112 kDa
Isotype: Observed MW:
IgG 66-75 kDa

Immunogen Catalog Number:

AG17480

Purification Method:

Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:50-1:500 IF 1:50-1:500

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, WB

IP: HepG2 cells,

Species Specificity: IHC : human testis tissue, human ovary tumor tissue

IF: Caco-2 cells, NCCIT cells

human, rat, mouse
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

SALL4, also named Sal-like protein 4 or Zinc finger protein 797, Contains 7 C2H2-type zinc fingers and belongs to the sal C2H2-type zinc-finger protein family. SALL4 is constitutively expressed in acute myeloid leukemia. The constitutive expression of SALL4 in mice is sufficient to induce MDS-like symptoms and transformation to AML that is transplantable. SALL4 is able to bind beta-catenin and activate the Wnt/beta-catenin signaling pathway. Sequence analysis of the larger cDNA fragment isolated revealed a single, large open-reading frame, designated as SALL4A, that started from a strong consensus initiation sequence and was expected to encode 1053 amino acids. The other splicing variant of SALL4, designated SALL4B, lacked the region corresponding to amino acids 385 to 820 of the full-length SALL4A. The putative protein encoded by SALL4B cDNA was expected to consist of 617 amino acids.

Notable Publications

Author	Pubmed ID	Journal	Application
Chaoqun Liu	34551797	J Exp Clin Cancer Res	WB
Qing-Dong Wang	36285444	Pathol Int	WB
Honghai Xia	27725724	Sci Rep	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




HepG2 cells were subjected to SDS PAGE followed by western blot with 24500-1-AP (SALL4 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human testis slide using 24500-1-AP (SALL4 Antibody) at dilution of 1:50.




Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using SALL4 antibody (24500-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using SALL4 antibody (24500-1-AP) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed NCCIT cells using SALL4 antibody (24500-1-AP) at dilution of 1:2000 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP Result of anti-SALL4 (IP:24500-1-AP, 4ug; Detection:24500-1-AP 1:600) with HepG2 cells lysate 3600ug.