For Research Use Only

## Angiopoietin 2 Polyclonal antibody Catalog Number: 24613-1-AP 29 Publications



Basic Information	Catalog Number: 24613-1-AP	GenBank Accession Number: BC126200	Purification Method: Antigen affinity purification
	Size:	GeneID (NCBI):	Recommended Dilutions:
	150ul , Concentration: 650 ug/ml by	285	WB: 1:500-1:2000
	Nanodrop and 367 ug/ml by Bradford	UNIPROT ID:	IF/ICC: 1:200-1:800
	method using BSA as the standard;	015123	
	Source: Rabbit	Full Name: angiopoietin 2	
	lsotype: IgG	Calculated MW: 496 aa, 57 kDa	
	Immunogen Catalog Number: AG17938	Observed MW: 68 kDa	
Applications	Tested Applications:	Positive C	Controls:
	WB, IF/ICC, ELISA	WB : HUVE	EC cells, HeLa cells, HepG2 cells, K-562 cells
	Cited Applications: WB, IHC, IF	IF/ICC : PO	C-12 cells, A549 cells
	Species Specificity: human		
	Cited Species: human, mouse, rat		
Background Informatior	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathw. identified through a cDNA library scre natural antagonist for Tie2 that disrup	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult n modeling. Angiopoietin-2 has also	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a nice and humans, Angiopoietin-2 is b been found to be highly expressed in
Background Informatior	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathw. identified through a cDNA library scre natural antagonist for Tie2 that disruj expressed only at sites of vascular re diverse tumor cells and plays an imp	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult n modeling. Angiopoietin-2 has also	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a nice and humans, Angiopoietin-2 is b been found to be highly expressed in
	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathw identified through a cDNA library scre natural antagonist for Tie2 that disrup expressed only at sites of vascular re diverse tumor cells and plays an imp	ay, one of the main pathways invo cening, shortly after the identificat pts in vivo angiogenesis. In adult n modeling. Angiopoietin-2 has also ortant role in tumor angiogenesis a	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a mice and humans, Angiopoietin-2 is been found to be highly expressed in and inflammation.
	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathw identified through a cDNA library scre natural antagonist for Tie2 that disrup expressed only at sites of vascular re diverse tumor cells and plays an imp Author Public Lifeng Wang 3610	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult m modeling. Angiopoietin-2 has also ortant role in tumor angiogenesis a med ID Journal	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a nice and humans, Angiopoietin-2 is been found to be highly expressed in and inflammation. Application
	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathwi identified through a cDNA library scree natural antagonist for Tie2 that disruj expressed only at sites of vascular re- diverse tumor cells and plays an imp Author Pub Lifeng Wang 3610 Qian Li 277	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult n modeling. Angiopoietin-2 has also ortant role in tumor angiogenesis a med ID Journal 60014 Front Genet	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a mice and humans, Angiopoietin-2 is been found to be highly expressed in and inflammation. Application IHC
	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathwi identified through a cDNA library scree natural antagonist for Tie2 that disruj expressed only at sites of vascular re- diverse tumor cells and plays an imp Author Pub Lifeng Wang 3610 Qian Li 277	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult m modeling. Angiopoietin-2 has also ortant role in tumor angiogenesis a med ID Journal 60014 Front Genet 71855 Tumour Biol	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a mice and humans, Angiopoietin-2 is been found to be highly expressed in and inflammation. Application IHC WB
Notable Publications	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathwi identified through a cDNA library scree natural antagonist for Tie2 that disruj expressed only at sites of vascular re- diverse tumor cells and plays an imp Author Pub Lifeng Wang 3610 Qian Li 277	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult m modeling. Angiopoietin-2 has also ortant role in tumor angiogenesis a med ID Journal 60014 Front Genet 71855 Tumour Biol 19207 Endocrinology er shipment.	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a mice and humans, Angiopoietin-2 is been found to be highly expressed in and inflammation. Application IHC WB
	human, mouse, rat Angiopoietin-2 (Ang2) is a growth fac homology domains) signaling pathwi identified through a cDNA library scree natural antagonist for Tie2 that disruj expressed only at sites of vascular re- diverse tumor cells and plays an imp Author Pub Lifeng Wang 3611 Qian Li 2777 Qiuyan Ma 3623 Storage: Store at -20°C. Stable for one year aft Storage Buffer:	ay, one of the main pathways invo eening, shortly after the identificat pts in vivo angiogenesis. In adult m modeling. Angiopoietin-2 has also ortant role in tumor angiogenesis a med ID Journal 60014 Front Genet 71855 Tumour Biol 19207 Endocrinology er shipment.	olved in angiogenesis. Angiopoietin-2 was tion of angiopoietin-1. Angiopoietin-2, is a mice and humans, Angiopoietin-2 is been found to be highly expressed in and inflammation. Application IHC WB

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Immunofluorescent analysis of (-20°C Methanol) fixed A549 cells using Angiopoietin 2 antibody (24613-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed PC-12 cells using Angiopoietin 2 antibody (24613-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Various lysates were subjected to SDS PAGE followed by western blot with 24613-1-AP (Angiopoietin 2 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.