For Research Use Only

PIK3C2B Polyclonal antibody

Catalog Number:24788-1-AP 2 Publications




Nam Sc Ra Is Ig An Ac An Ci W Sp hu Ci Notable Publications	Oul, Concentration: 450 ug/ml by nodrop and 367 ug/ml by Bradfor ethod using BSA as the standard; urce: bbit	d UNIPROT IE 000750 Full Name: phosphoinc beta polype Calculated 1634 aa, 18 Observed N	D: Disitide-3-kinase, class 2, eptide MW:	Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:100-1:400 IF/ICC 1:10-1:100	
Advantage Advant		Observed N	35 kDa	WB 1:1000-1:6000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:100-1:400	
Notable Publications		1634 aa, 185 kDa Observed MW: 160-185 kDa			
Ci W Sp hu Ci hu N Ti re bu Notable Publications	sted Applications: 3, IHC, IF/ICC, IP, ELISA		Positive Controls:		
Sp hu Ci hu N Ti re bu Notable Publications	Cited Applications:		WB : HeLa cells, Jurkat cells IP : HeLa cells,		
Notable Publications	WB Species Specificity:		IHC : human skeletal muscle tissue,		
hu N Ti re bu Notable Publications	man		IF/ICC : HeLa	cells,	
Notable Publications	red Species: man				
Jia	ote-IHC: suggested antigen buffer pH 9.0; (*) Alternat trieval may be performed v ffer pH 6.0	ively, antig	en		
	thor Pu	bmed ID	Journal	Application	
RL	wen Lyu 29	660285	Anal Chem		
	-Xu Sun 37	436898	Cell Rep	WB	
Storage St St PE	orage: ore at -20°C. Stable for one year a orage Buffer: S with 0.02% sodium azide and 5	0% glycerol p			
Al *** 20ul sizes contain 0.1% BSA		storage			


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 24788-1-AP (PIK3C2B antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




IP result of anti-PIK3C2B (IP:24788-1-AP, 4ug; Detection:24788-1-AP 1:1500) with HeLa cells lysate 1240 ug.



Immunohistochemical analysis of paraffinembedded human skeletal muscle tissue slide using 24788-1-AP (PIK3C2B Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human skeletal muscle tissue slide using 24788-1-AP (PIK3C2B Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using 24788-1-AP (PIK3C2B antibody) at dilution of 1:25 and Alexa Fluor 488conjugated Goat Anti-Rabbit IgG(H+L).