

For Research Use Only

TNRC4 Polyclonal antibody

Catalog Number: 25353-1-AP



Basic Information

Catalog Number: 25353-1-AP	GenBank Accession Number: BC052491	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 550 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 11189	Recommended Dilutions: WB 1:500-1:2000 IHC 1:20-1:200
Source: Rabbit	Full Name: trinucleotide repeat containing 4	
Isotype: IgG	Calculated MW: 464 aa, 51 kDa	
Immunogen Catalog Number: AG18601	Observed MW: 55 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse lung tissue, mouse spleen tissue

IHC : human testis tissue, human cerebellum tissue

Background Information

TNRC4, also named as BRUNOL1, CAGH4, ERDA4, or TNRC4, is a 465 amino acid protein, which contains three RRM (RNA recognition motif) domains and belongs to the CELF/BRUNOL family. TNRC4 localizes in the nucleus and is expressed in brain. TNRC4 as a RNA-binding protein is involved in the regulation of pre-mRNA alternative splicing and mediates exon inclusion and/or exclusion in pre-mRNA that are subject to tissue-specific and developmentally regulated alternative splicing.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



mouse lung tissue were subjected to SDS PAGE followed by western blot with 25353-1-AP (TNRC4 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.

