For Research Use Only

TTC35 Polyclonal antibody

Catalog Number: 25443-1-AP

Featured Product

13 Publications



Basic Information

Catalog Number: GenBank Accession Number:

25443-1-AP BC021667 GeneID (NCBI): Size:

150ul, Concentration: 1200 ug/ml by 9694 Nanodrop and 507 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; Q15006

Source: Full Name:

Rabbit tetratricopeptide repeat domain 35

Isotype: Calculated MW: 297 aa, 35 kDa Immunogen Catalog Number: Observed MW: AG22196 35 kDa

Purification Method: Antigen affinity purification Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

WB 1:1000-1:8000 protein lysate

Applications

Tested Applications: WB, IP, ELISA

Cited Applications:

WB, IP, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse

Positive Controls:

WB: Jurkat cells, mouse heart tissue, K-562 cells, MCF-7 cells, NIH/3T3 cells, mouse spleen tissue, human placenta tissue, PC-3 cells, SKOV-3 cells

IP: mouse heart tissue,

Background Information

TPR repeat protein 35 (TTC35), also named as tetratricopeptide repeat domain 35, is a 297 amino acid protein, which contains three TPR repeats and belongs to the EMC2 family. TTC35 localizes in the nucleus and cytoplasm. TTC35 is a component of the ER membrane protein complex and may be a putative O-linked glycosyl transferase.

Notable Publications

Author	Pubmed ID	Journal	Application
Ashley M Ngo	31516121	Elife	WB
Patrick J Chitwood	30415835	Cell	WB
Cary R Boyd-Shiwarski	36318922	Cell	IF

Storage

Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 25443-1-AP (TTC35 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



IP result of anti-TTC35 (IP:25443-1-AP, 3ug; Detection:25443-1-AP 1:600) with mouse heart tissue lysate 4000ug.



Various lysates were subjected to SDS PAGE followed by western blot with 25443-1-AP (TTC35 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.