For Research Use Only

c-Cbl Polyclonal antibody

Catalog Number:25818-1-AP

Featured Product

19 Publications



Basic Information	Catalog Number: 25818-1-AP	GenBank Accession Number: BC 132733	Purification Method: Antigen affinity purification	
	Size:	GenelD (NCBI):	Recommended Dilutions:	
	150ul , Concentration: 600 ug/ml by	867	WB 1:5000-1:50000	
	Nanodrop;	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total	
	Source:	P22681	protein lysate IHC 1:50-1:500	
	Rabbit	Full Name:		
	lsotype: IgG	Cas-Br-M (murine) ecotropic retroviral transforming sequence		
	Immunogen Catalog Number: AG23033	Calculated MW: 906 aa, 100 kDa		
		Observed MW: 100-120 kDa		
Applications	Tested Applications: WB, IHC, IP, ELISA	Positive Controls:		
	Cited Applications:	WB : K-562 cells, Raji cells, Ramos cells, Romas cells		
	WB, IHC, IF, IP, CoIP	IP : K-562 cells,		
	Species Specificity: human, mouse	Inc : numa	an tonsillitis tissue, human lymphoma tissue	
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	TE buffer pH 9.0; (*) Alternati retrieval may be performed w	ively, antigen		
Background Information	TE buffer pH 9.0; (*) Alternati retrieval may be performed v buffer pH 6.0 c-Cbl (Casitas B-lineage lymphoma) development, B cell development, in	ively, antigen with citrate I is a critical regulator that controls i mmune tolerance, angiogenesis, or ndritic cell (DC) function via negativ	immature or follicular helper T cell tumorigenesis (PMID: 33962939). c-Cbl also vely modulating IL-12 production after Toll-	
	TE buffer pH 9.0; (*) Alternati retrieval may be performed v buffer pH 6.0 c-Cbl (Casitas B-lineage lymphoma) development, B cell development, in plays a pivotal role in regulating de like receptor (TLR) stimulation (PMIC	ively, antigen with citrate I is a critical regulator that controls i mmune tolerance, angiogenesis, or ndritic cell (DC) function via negativ	tumorigenesis (PMID: 33962939). c-Cbl also	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed we buffer pH 6.0 c-Cbl (Casitas B-lineage lymphoma) development, B cell development, in plays a pivotal role in regulating de like receptor (TLR) stimulation (PMID Author Pul	ively, antigen with citrate I is a critical regulator that controls in mmune tolerance, angiogenesis, or ndritic cell (DC) function via negativ D: 21799517).	tumorigenesis (PMID: 33962939). c-Cbl also vely modulating IL-12 production after Toll-	
Background Information Notable Publications	TE buffer pH 9.0; (*) Alternative retrieval may be performed weight buffer pH 6.0 c-Cbl (Casitas B-lineage lymphoma) development, B cell development, in plays a pivotal role in regulating de like receptor (TLR) stimulation (PMIC Author Pul Jennifer J Schwarz 343	ively, antigen with citrate his a critical regulator that controls in mmune tolerance, angiogenesis, or ndritic cell (DC) function via negative D: 21799517).	tumorigenesis (PMID: 33962939). c-Cbl also vely modulating IL-12 production after Toll- Application WB	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed web buffer pH 6.0 c-Cbl (Casitas B-lineage lymphoma) development, B cell development, in plays a pivotal role in regulating de like receptor (TLR) stimulation (PMID Author Pul Jennifer J Schwarz 342 Yan Yang 362	ively, antigen with citrate It is a critical regulator that controls in mmune tolerance, angiogenesis, or indritic cell (DC) function via negative D: 21799517). Demed ID Journal 526379 Life Sci Alliance	tumorigenesis (PMID: 33962939). c-Cbl also vely modulating IL-12 production after Toll- Application WB	
	TE buffer pH 9.0; (*) Alternative retrieval may be performed web buffer pH 6.0 c-Cbl (Casitas B-lineage lymphoma) development, B cell development, in plays a pivotal role in regulating de like receptor (TLR) stimulation (PMID Author Pul Jennifer J Schwarz 342 Yan Yang 362	ively, antigen with citrate	tumorigenesis (PMID: 33962939). c-Cbl also vely modulating IL-12 production after Toll- Application WB col WB,IP	


For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 25818-1-AP (c-Cbl antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 25818-1-AP (c-Cbl antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.



IP result of anti-c-Cbl (IP:25818-1-AP, 4ug; Detection:25818-1-AP 1:1000) with K-562 cells lysate 3680 ug.



Immunohistochemical analysis of paraffinembedded human lymphoma tissue slide using 25818-1-AP (c-Cbl antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).