For Research Use Only

KIF5C Polyclonal antibody Catalog Number:25897-1-AP Featured Product

Featured Product

9 Publications



Basic Information	Catalog Number: 25897-1-AP	GenBank Accession Number: BC110287	Purification Method: Antigen affinity purification
	Size:	GenelD (NCBI):	Recommended Dilutions:
	150ul , Concentration: 1500 ug/ml by	3800	WB 1:1000-1:8000
	Nanodrop and 787 ug/ml by Bradford	UNIPROT ID:	IP 0.5-4.0 ug for 1.0-3.0 mg of total
	method using BSA as the standard;	060282	protein lysate IHC 1:50-1:500
	Source: Rabbit	Full Name:	IF/ICC 1:50-1:500
	Isotype:	kinesin family member 5C	
	lgG	Calculated MW: 725 aa, 83 kDa	
	Immunogen Catalog Number: AG23163	Observed MW: 70 kDa, 110 kDa	
Applications			re Controls:
	Cited Applications:		ouse brain tissue, rat brain tissue
	WB, IHC, CoIP		use brain tissue,
	Species Specificity: human, mouse, rat		nouse brain tissue, : HeLa cells, SH-SY5Y cells
	Cited Species:	in/rec	
	human, mouse		
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ		
	retrieval may be performed w buffer pH 6.0	ith citrate	
Background Information	buffer pH 6.0 KIF5C (Kinesin heavy chain isoform 5 The kinesin-1 subfamilies (kif5a, kif5 Kinesins have N-terminal motor dom The hinge and C-terminal tail regions granule. The granule localized to den the granule was enhanced by Kif5 over	C variant) which belongs to the b, kif5c) are responsible for the ains and C-terminal cargo-bind s of Kif5a, Kif5b, and Kif5c bound drites and underwent bidirection erexpression and reduced by Kif e. Two major isoforms of kif5c,	e movement of mitochondria in neurons. ling tail domains separated by hinge regions. d a large detergent-resistant RNase-sensitive onal movement. Distally directed movement of f5 functional blockage. kinesins may transpor
	buffer pH 6.0 KIF5C (Kinesin heavy chain isoform 5 The kinesin-1 subfamilies (kif5a, kif5 Kinesins have N-terminal motor dom The hinge and C-terminal tail regions granule. The granule localized to den the granule was enhanced by Kif5 ove RNA in dendrites via this large granul kDa, are generated by alternative spl	C variant) which belongs to the b, kif5c) are responsible for the ains and C-terminal cargo-bind s of Kif5a, Kif5b, and Kif5c bound drites and underwent bidirection erexpression and reduced by Kif e. Two major isoforms of kif5c,	ing tail domains separated by hinge regions.
	buffer pH 6.0 KIF5C (Kinesin heavy chain isoform 5 The kinesin-1 subfamilies (kif5a, kif9 Kinesins have N-terminal motor dom The hinge and C-terminal tail regions granule. The granule localized to den the granule was enhanced by Kif5 ove RNA in dendrites via this large granul kDa, are generated by alternative spl Author Pub	C variant) which belongs to the 5b, kif5c) are responsible for the ains and C-terminal cargo-bind s of Kif5a, Kif5b, and Kif5c bound drites and underwent bidirectic arexpression and reduced by Kit e. Two major isoforms of kif5c, icing.	e movement of mitochondria in neurons. ling tail domains separated by hinge regions. d a large detergent-resistant RNase-sensitive onal movement. Distally directed movement of f5 functional blockage. kinesins may transpor with molecular predicted masses of 80 and 13 Application
Background Information	buffer pH 6.0 KIF5C (Kinesin heavy chain isoform 5 The kinesin-1 subfamilies (kif5a, kif5 Kinesins have N-terminal motor dom. The hinge and C-terminal tail regions granule. The granule localized to den the granule was enhanced by Kif5 over RNA in dendrites via this large granul kDa, are generated by alternative spl Author Pub Tien-Chun Yang 322	C variant) which belongs to the 5b, kif5c) are responsible for the ains and C-terminal cargo-bind s of Kif5a, Kif5b, and Kif5c bound drites and underwent bidirectic erexpression and reduced by Kii e. Two major isoforms of kif5c, icing. med ID Journal	e movement of mitochondria in neurons. ling tail domains separated by hinge regions. d a large detergent-resistant RNase-sensitive onal movement. Distally directed movement of f5 functional blockage. kinesins may transpor with molecular predicted masses of 80 and 13 Application
	buffer pH 6.0KIF5C (Kinesin heavy chain isoform 5The kinesin-1 subfamilies (kif5a, kif5Kinesins have N-terminal motor domThe hinge and C-terminal tail regionsgranule. The granule localized to denthe granule was enhanced by Kif5 overRNA in dendrites via this large granulkDa, are generated by alternative spleAuthorPubTien-Chun Yang322Qi Wang309	C variant) which belongs to the 5b, kif5c) are responsible for the ains and C-terminal cargo-bind s of Kif5a, Kif5b, and Kif5c bound drites and underwent bidirectic erexpression and reduced by Kit e. Two major isoforms of kif5c, icing. med ID Journal 77753 Hum Mol Gene	e movement of mitochondria in neurons. ling tail domains separated by hinge regions. d a large detergent-resistant RNase-sensitive onal movement. Distally directed movement of f5 functional blockage. kinesins may transpor with molecular predicted masses of 80 and 1 Application et WB WB
	buffer pH 6.0KIF5C (Kinesin heavy chain isoform 5The kinesin-1 subfamilies (kif5a, kif5Kinesins have N-terminal motor domThe hinge and C-terminal tail regionsgranule. The granule localized to denthe granule was enhanced by Kif5 overRNA in dendrites via this large granulkDa, are generated by alternative spleAuthorPubTien-Chun Yang322Qi Wang309	iC variant) which belongs to the 5b, kif5c) are responsible for the ains and C-terminal cargo-bind s of Kif5a, Kif5b, and Kif5c bound drites and underwent bidirectic erexpression and reduced by Kit e. Two major isoforms of kif5c, icing. med ID Journal 77753 Hum Mol Gene 23004 Neurobiol Dis 91783 Alzheimers Re er shipment. % glycerol pH 7.3.	e movement of mitochondria in neurons. ling tail domains separated by hinge regions. d a large detergent-resistant RNase-sensitive onal movement. Distally directed movement of f5 functional blockage. kinesins may transpor with molecular predicted masses of 80 and 1 Application et WB WB


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com

in USA), or 1(312) 455-8498 (outside USA)


W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 25897-1-AP (KIF5C antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




IP result of anti-KIF5C (IP:25897-1-AP, 4ug; Detection:25897-1-AP 1:800) with mouse brain tissue lysate 2080 ug.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 25897-1-AP (KIF5C antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using KIF5C antibody (25897-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using KIF5C antibody (25897-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).