#### For Research Use Only

# **BDNF Polyclonal antibody**

Catalog Number: 28205-1-AP 152 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

28205-1-AP BC029795
Size: GeneID (NCBI):

150ul , Concentration: 700 ug/ml by 627
Nanodrop; UNIPROT ID:
Source: P23560
Rabbit Full Name:

Isotype: brain-derived neurotrophic factor

IgG Calculated MW:
Immunogen Catalog Number: 247 aa, 28 kDa
AG28276 Observed MW:
28-30 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB: 1:1000-1:8000 IHC: 1:400-1:1600

**Applications** 

Tested Applications: WB, IHC, ELISA

Cited Applications: WB, IHC, IF

Species Specificity: human, mouse, rat Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: mouse brain tissue, mouse cerebellum tissue, PC-12 cells, SH-SY5Y cells, rat brain tissue, rat cerebellum, C6 cells

IHC: mouse cerebellum tissue,

## **Background Information**

BDNF is a member of the family of neurotrophic factors. It was first explored and purified from pig brain in 1982, and other members of the neurotrophin family were successively discovered. BDNF is induced by cortical neurons, and is necessary for survival of striatal neurons in the brain. Expression of BDNF is reduced in both Alzheimer's and Huntington disease patients. It participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. It also plays a role in the regulation of stress response and in the biology of mood disorders. The BDNF gene is rather complex in its structure and it can express five different mRNA isoforms by alternative splicing, producing 28-37 kDa proteins.

#### Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoyan Liu	31574948	Int J Mol Sci	IF
Minghao Yao	31355388	Biomater Sci	WB
Weiping Xiao	36100045	Exp Neurol	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


### **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 28205-1-AP (BDNF antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 28205-1-AP (BDNF antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 28205-1-AP (BDNF antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).