

For Research Use Only

PSMB11 Polyclonal antibody

Catalog Number: 55143-1-AP



Basic Information

Catalog Number: 55143-1-AP	GenBank Accession Number: NM_001099780	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 122706	Recommended Dilutions: WB 1:200-1:600 IHC 1:100-1:400
Source: Rabbit	Full Name: proteasome (prosome, macropain) subunit, beta type, 11	
Isotype: IgG	Calculated MW: 33 kDa	
	Observed MW: 32-35 kDa	

Applications

Tested Applications: FC, IHC, WB, ELISA	Positive Controls:
Species Specificity: human	WB : Jurkat cells, IHC : human thymus tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PSMB11 belongs to the peptidase T1B family. It is a multicatalytic proteinase complex which is characterized by its ability to cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. PSMB11 has an ATP-dependent proteolytic activity. Incorporated instead of PSMB5 or PSMB8, this unit reduces the chymotrypsin-like activity of the proteasome. It plays a pivotal role in development of CD8-positive T cells. The antibody is specific to PSMB11.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 55143-1-AP (PSMB11 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human thymus tissue slide using 55143-1-AP (PSMB11 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human thymus tissue slide using 55143-1-AP (PSMB11 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1×10^6 MCF-7 cells were intracellularly stained with 0.2 ug Anti-Human PSMB11 (55143-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).