For Research Use Only

STUB1 Polyclonal antibody

Catalog Number: 55430-1-AP

Featured Product 33 Publications



Basic Information	Catalog Number: 55430-1-AP	GenBank Accession Numb NM_005861	er: Purification Method: Antigen affinity purification
	Size: 150ul , Concentration: 500 ug/ml by Nanodrop;	GenelD (NCBI): 10273	Recommended Dilutions: WB: 1:500-1:1000 IHC: 1:50-1:500
	Source:	UNIPROT ID: Q9UNE7	mc. 1.90 1.900
	Rabbit	Full Name:	
	Isotype:STIP1 homologyIgGcontaining prote		х
		Calculated MW: 35 kDa	
		Observed MW: 35 kDa	
Applications	Tested Applications:	Positive Controls:	
	WB, IHC, ELISA Cited Applications:	WB : HeLa cells, mouse brain tissue, HEK-293 cells, MCF-7 cells, mouse liver tissue	
	WB, IHC, IF, IP Species Specificity: human, mouse, rat	becificity: mouse skeletal muscle tissue, human heart t	
	Cited Species: human, mouse, rat	nu	
	Note-IHC: suggested antigen r TE buffer pH 9.0; (*) Alternativ retrieval may be performed w buffer pH 6.0	vely, antigen	
	STUB1, also named as CHIP, SDCCAG7, UBOX1 and NY-CO-7, is a E3 ubiquitin-protein ligase which targets misfolded chaperone substrates towards proteasomal degradation. STUB1/CHIP modulates the activity of several chaperone complexes, including Hsp70, Hsc70 and Hsp90. It mediates transfer of non-canonical short ubiquitin chains to HSPA8 that have no effect on HSPA8 degradation. STUB1/CHIP mediates polyubiquitination of CYP3A4. (PMID:10330192) CHIP/STUB1 strongly reduced keratin aggregates through increased degradation of mutant K14. (PMID:20151404) STUB1/CHIP regulates NAD(P)H:quinone oxidoreductase 1 (NQO1) accumulation in aged brain, a process impaired in certain Alzheimer patients.(PMID:21220432) This antibody is specific to STUB1.		
Background Information	misfolded chaperone substrates towa chaperone complexes, including Hsp chains to HSPA8 that have no effect of (PMID:10330192) CHIP/STUB1 strong (PMID:20151404) STUB1/CHIP regula	ards proteasomal degradati 70, Hsc70 and Hsp90. It med on HSPA8 degradation. STUB ly reduced keratin aggrega tes NAD(P)H:quinone oxido	on. STUB1/CHIP modulates the activity of severa diates transfer of non-canonical short ubiquitin B1/CHIP mediates polyubiquitination of CYP3A4 tes through increased degradation of mutant K14 reductase 1 (NQO1) accumulation in aged brain,
	misfolded chaperone substrates towa chaperone complexes, including Hsp chains to HSPA8 that have no effect of (PMID:10330192) CHIP/STUB1 strong (PMID:20151404) STUB1/CHIP regula process impaired in certain Alzheime	ards proteasomal degradati 70, Hsc70 and Hsp90. It med on HSPA8 degradation. STUB ly reduced keratin aggrega tes NAD(P)H:quinone oxido	on. STUB1/CHIP modulates the activity of severa diates transfer of non-canonical short ubiquitin B1/CHIP mediates polyubiquitination of CYP3A4. tes through increased degradation of mutant K14 reductase 1 (NQO1) accumulation in aged brain,
	misfolded chaperone substrates towa chaperone complexes, including Hsp chains to HSPA8 that have no effect of (PMID:10330192) CHIP/STUB1 strong (PMID:20151404) STUB1/CHIP regula process impaired in certain Alzheime Author Put	ards proteasomal degradati 70, Hsc70 and Hsp90. It med In HSPA8 degradation. STUE ly reduced keratin aggrega tes NAD(P)H:quinone oxidd er patients.(PMID:21220432	on. STUB1/CHIP modulates the activity of severa liates transfer of non-canonical short ubiquitin 81/CHIP mediates polyubiquitination of CYP3A4 tes through increased degradation of mutant K14 reductase 1 (NQO1) accumulation in aged brain,) This antibody is specific to STUB1. Application
	misfolded chaperone substrates towa chaperone complexes, including Hsp chains to HSPA8 that have no effect of (PMID:10330192) CHIP/STUB1 strong (PMID:20151404) STUB1/CHIP regula process impaired in certain Alzheime Author Put Hao Chen 330	ands proteasomal degradati 70, Hsc70 and Hsp90. It med on HSPA8 degradation. STUE ly reduced keratin aggrega tes NAD(P)H:quinone oxido er patients.(PMID:21220432 med ID Journal	on. STUB1/CHIP modulates the activity of severa liates transfer of non-canonical short ubiquitin 81/CHIP mediates polyubiquitination of CYP3A4 tes through increased degradation of mutant K14 reductase 1 (NQO1) accumulation in aged brain,) This antibody is specific to STUB1. Application
	misfolded chaperone substrates towa chaperone complexes, including Hsp chains to HSPA8 that have no effect of (PMID:10330192) CHIP/STUB1 strong (PMID:20151404) STUB1/CHIP regula process impaired in certain Alzheime Author Put Hao Chen 330 Sheng Li 347	ards proteasomal degradati 70, Hsc70 and Hsp90. It med on HSPA8 degradation. STUE ly reduced keratin aggrega tes NAD(P)H:quinone oxido er patients.(PMID:21220432 med ID Journal 013306 Front Neur	on. STUB1/CHIP modulates the activity of severa liates transfer of non-canonical short ubiquitin B1/CHIP mediates polyubiquitination of CYP3A4 tes through increased degradation of mutant K14 reductase 1 (NQO1) accumulation in aged brain,) This antibody is specific to STUB1. Application rosci WB WB
Background Information Notable Publications	misfolded chaperone substrates towa chaperone complexes, including Hsp chains to HSPA8 that have no effect of (PMID:10330192) CHIP/STUB1 strong (PMID:20151404) STUB1/CHIP regula process impaired in certain Alzheime Author Put Hao Chen 330 Sheng Li 347	ards proteasomal degradati 70, Hsc70 and Hsp90. It med on HSPA8 degradation. STUE ly reduced keratin aggrega tes NAD(P)H:quinone oxidd er patients.(PMID:21220432 med ID Journal 013306 Front Neu 01988 Nat Comm er shipment. % glycerol, pH7.3	on. STUB1/CHIP modulates the activity of severa liates transfer of non-canonical short ubiquitin B1/CHIP mediates polyubiquitination of CYP3A4 tes through increased degradation of mutant K14 reductase 1 (NQO1) accumulation in aged brain,) This antibody is specific to STUB1. Application rosci WB WB

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com

in USA), or 1(312) 455-8498 (outside USA)


W: ptglab.com

Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 55430-1-AP (STUB1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of STUB1 antibody (55430-1-AP, 1:500) with si-control and si-STUB1 transfected HeLa cells.