For Research Use Only

BRD8 Monoclonal antibody

Catalog Number:60121-1-lg Featured Product




Basic Information	Catalog Number: 60121-1-lg	GenBank Accession Number: BC008039 GeneID (NCBI): 10902		Purification Method: Protein A purification CloneNo.: 5B6D6	
	150ul, Concentration: 1467 ug/ml by Bradford method using BSA as the standard; Source:				
		UNIPROT ID: Q9H0E9 Full Name: bromodomain containing 8 Calculated MW: 120 kDa		Recommended Dilutions: WB 1:500-1:2000 IHC 1:50-1:500 IF/ICC 1:1000-1:4000	
					lsotype: lgG1
	Immunogen Catalog Number: AG0790				Observed MW: 120 kDa
	Applications	Tested Applications:	Positive Co		rols:
WB, IHC, IF/ICC, ELISA		٧	WB: fetal human brain tissue, HeLa cells		
Species Specificity: human, mouse, rat		I	IHC : human colon cancer tissue,		
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		F/ICC : HepG	2 cells, HEK-293 cells		
Background Information	BRD8 is one of the bromodomain-containing proteins, which is an acetylated lysine- binding domain and thought to be involved in regulation of protein acetylation and/or HAT (histone acetyl transferase) activity [PMID: 12963728]. It is also reported to be an accessory subunit of NuA4 HAT complex (also known as TRRAP/TIP60 complex) through biochemical purification [PMID: 16049979]. In addition, it can bind to thyroid hormone receptor-ß and/or retinoid X receptor, and can act as a coactivator of the nuclear hormone receptor-mediated transcription in reporter assays [PMID: 10517671].				
	Storage: Store at -20°C. Stable for one year aft	er shinment			
Storage	Storage Buffer: PBS with 0.02% sodium azide and 50				

For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of BRD8 antibody (60121-1-lg, 1:8000) with si-Control and si-BRD8 transfected HeLa cells..

250kd→ 150kd→ 100kd→ 70kd→ 50kd→ 40kd→ 30kd→

fetal human brain tissue were subjected to SDS PAGE followed by western blot with 60121-1-1g (BRD8 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 60121-1-1g (BRD8 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using BRD8 antibody (60121-1-1g, Clone: 5B6D6) at dilution of 1:2000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).