For Research Use Only

FGF12 Monoclonal antibody

Catalog Number:60152-1-lg





Basic Information	Catalog Number: 60152-1-lg	GenBank Accession Number: BC022524	Purification Method: Protein A purification	
	Size: 150ul , Concentration: 1200 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2257	CloneNo.: 2F11A10	
		UNIPROT ID: P61328	Recommended Dilutions: WB 1:2000-1:10000	
	Source: Mouse	Full Name: fibroblast growth factor 12		
	Isotype: Calculated MW: IgG2b 181 aa, 20 kDa			
	Immunogen Catalog Number: AG5046	Observed MW: 20 kDa		
Applications	Tested Applications: WB, ELISA, FC (Intra)		ositive Controls:	
	Species Specificity: tissue human tissue		use brain tissue, human brain tissue, rat brain	
Background Information	FGF12 is a member of the fibroblast growth factor (FGF) family. FGF family members play important roles in embryogenesis, angiogenesis, and wound repair. FGF12 lacks the N-terminal signal sequence present in most of the FGF family members, but it contains clusters of basic residues that have been demonstrated to act as a nuclear localization signal. When transfected into mammalian cells, this protein accumulated in the nucleus, but was not secreted. FGF12 plays an intracellular role in the inhibition of radiation-induced apoptosis. FGF12 is expressed abundantly in developing and adult nervous systems; therefore, FGF12 was thought to be related to nervous system development and function.			
Storage	Storage: Store at -20°C. Stable for one year after Storage Buffer: PBS with 0.02% sodium azide and 50°			
	Aliquoting is unnecessary for -20° C s	torage		

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)E: proteintech@ptglab.comW: ptglab.comW: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse brain tissue were subjected to SDS PAGE followed by western blot with 60152-1-1g (FGF12 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. 1x10^6 HeLa cells were intracellularly stained with 0.8 ug FGF12 Monoclonal antibody (60152-1-1g, Clone:2F11A10) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse 1gG(H+L) (SA00013-1) (red), or 0.8 ug Mouse 1gG2b isotype control Mouse McAb (66360-3-1g, Clone: 11B8C4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).