

NCAM1/CD56 Monoclonal antibody

Catalog Number: 60238-1-Ig **9 Publications**

Basic Information

Catalog Number: 60238-1-Ig	GenBank Accession Number: BC047244	Purification Method: Protein G purification
Size: 150ul, Concentration: 1500 ug/ml by Nanodrop and 839 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 4684	CloneNo.: 1E8C9
Source: Mouse	ENSEMBL Gene ID: ENSG00000149294	Recommended Dilutions: WB: 1:1000-1:8000 IHC: 1:2000-1:8000 IF-P: 1:200-1:800 IF/ICC: 1:200-1:800
Isotype: IgG1	UNIPROT ID: P13591	
Immunogen Catalog Number: AG5732	Full Name: neural cell adhesion molecule 1	
	Calculated MW: 95 kDa	
	Observed MW: 140 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, IF-P, ELISA	Positive Controls:
Cited Applications: WB, IHC, IF	WB : fetal human brain tissue, C6 cells, human brain tissue, pig brain tissue
Species Specificity: human, rat, pig	IHC : human lung cancer tissue, human colon tissue
Cited Species: human, mouse	IF-P : human colon tissue, human tonsillitis tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IF/ICC : SH-SY5Y cells, human colon tissue

Background Information

Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronectin-type III repeats (FNIII). All three forms can be posttranslationally modified by addition of polysialic acid (PSA) (PMID: 14976519). Several other isoforms have also been described (PMID: 1856291).

Notable Publications

Author	Pubmed ID	Journal	Application
Liang Gong	34761524	Small	IHC
Yan Huang	35578896	Jpn J Clin Oncol	IHC
Xianzhi Liu	35849032	Hepatology	IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




fetal human brain tissue were subjected to SDS PAGE followed by western blot with 60238-1-Ig (NCAM1/CD56 antibody at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 60238-1-Ig (NCAM1/CD56 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 60238-1-Ig (NCAM1/CD56 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human colon tissue using NCAM1/CD56 antibody (60238-1-Ig, Clone: 1E8C9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using NCAM1/CD56 antibody (60238-1-Ig, Clone: 1E8C9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 60238-1-Ig (NCAM1/CD56 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human tonsillitis tissue using NCAM1/CD56 antibody (60238-1-Ig, Clone: 1E8C9) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).