For Research Use Only

Anti-Mouse PD-1/CD279 (RMP1-30)

Catalog Number:65142-1-Ig



Basic Information

Catalog Number:

GenBank Accession Number: BC119179

programmed cell death 1

Purification Method: Protein A purification

65142-1-lg Size:

GeneID (NCBI):

CloneNo.:

100ug , 0.5 mg/ml

18566

RMP1-30

Source:

UNIPROT ID: Q02242 Recommended Dilutions:

Isotype:

Full Name:

FC: 0.5 ug per 10^6 cells in 100 μl

IgG2b, kappa

suspension

Applications

Tested Applications:

FC

rositive controts.

Species Specificity:

mouse

FC: Con-A stimulated BALB/c mouse splenocytes,

Background Information

Programmed cell death 1 (PD-1, also known as CD279) is an immunoinhibitory receptor that belongs to the CD28/CTLA-4 subfamily of the Ig superfamily. It is a 288 amino acid (aa) type I transmembrane protein composed of one Ig superfamily domain, a stalk, a transmembrane domain, and an intracellular domain containing an immunoreceptor tyrosine-based inhibitory motif (ITIM) as well as an immunoreceptor tyrosine-based switch motif (ITSM) (PMID: 18173375). PD-1 is expressed during thymic development and is induced in a variety of hematopoietic cells in the periphery by antigen receptor signaling and cytokines (PMID: 20636820). Engagement of PD-1 by its ligands PD-L1 or PD-L2 transduces a signal that inhibits T-cell proliferation, cytokine production, and cytokytic function (PMID: 19426218). It is critical for the regulation of T cell function during immunity and tolerance. Blockade of PD-1 can overcome immune resistance and also has been shown to have antitumor activity (PMID: 22658127; 23169436).

Storage


Storage:

Store at 2-8°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.09% sodium azide, pH7.3

Selected Validation Data



1X10^6 Con-A stimulated BALB/c mouse splenocytes were surface stained with 0.5 ug Anti-Mouse PD-1/CD279 (65142-1-Ig, Clone: RMP1-30) or 0.5 ug rat IgG2b isotype control, and FITC anti-rat IgG2b Antibody at dilution 1:100. Cells were not fixed.