For Research Use Only

CD31 Monoclonal antibody

Catalog Number: 66065-1-lg 28 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

66065-1-lg

GeneID (NCBI):

CloneNo.: 2A1E2

150ul, Concentration: 913 µg/ml by 5175 Bradford method using BSA as the

Immunogen Catalog Number:

BC022512

Recommended Dilutions:

standard;

ENSEMBL Gene ID: ENSG00000261371

WB 1:2000-1:8000

IHC 1:400-1:4000

Source: Mouse

Isotype:

AG1787

lgG2b

UNIPROT ID: P16284 Full Name:

platelet/endothelial cell adhesion

molecule

Calculated MW:

83 kDa

Observed MW:

120-130 kDa

Applications

Tested Applications:

WB, IF, IHC, ELISA

Cited Applications:

WB, IF, IHC

Species Specificity:

human

Cited Species:

Positive Controls:

WB: Jurkat cells, THP-1 cells, human placenta tissue

IHC: human liver cancer tissue, human tonsillitis

tissue, human breast cancer tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Platelet endothelial cell adhesion molecule-1 (PECAM-1, CD31) is a member of the immunoglobulin gene superfamily of cell adhesion molecules. CD31 is a transmembrane glycoprotein that is highly expressed on the surface of the endothelium, making up a large portion of its intracellular junctions. PECAM-1 is also present on the surface of hematopoietic cells and immune cells including platelets, monocytes, neutrophils, natural killer cells, megakaryocytes and some types of T-cell (PMID: 9011572). As well as its role in cell-cell adhesion, PECAM-1 functions as a signaling receptor, and is involved in important physiological events such as nitric oxide production, regulation of T-cell immunity and tolerance, leukocyte transendothelial migration and inflammation and angiogenesis (PMID: 21183735; 20978210; 17872453; 20634489).

Notable Publications

Author	Pubmed ID	Journal	Application
Shiro Matsubara	31533865	J Immunother Cancer	IF
Alina Jankowska-Konsur	27630298	Anticancer Res	
Miranda R Sun	33117819	Front Cell Dev Biol	IF

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Untreated and PNGase F-treated lysates of Jurkat cells were subjected to SDS PAGE followed by western blot with 66065-1-Ig (CD31 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).




Various lysates were subjected to SDS PAGE followed by western blot with 66065-1-1g (CD31 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66065-1-lg (CD31 antibody) at dilution of 1:4000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66065-1-Ig (CD31 antibody) at dilution of 1:4000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 66065-1-1g(CD31 antibody) at dilution of 1:200 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using 66065-1-1g(CD31 antibody) at dilution of 1:200 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).