

For Research Use Only

GM2A Monoclonal antibody

Catalog Number: 66080-1-Ig **Featured Product**



Basic Information

Catalog Number: 66080-1-Ig	GenBank Accession Number: BC009273	Purification Method: Protein A purification
Size: 150ul , Concentration: 2100 µg/ml by Nanodrop and 1640 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2760	CloneNo.: 1E4D3
Source: Mouse	Full Name: GM2 ganglioside activator	Recommended Dilutions: WB 1:500-1:1000
Isotype: IgG2a	Calculated MW: 22 kDa	
Immunogen Catalog Number: AG4394	Observed MW: 22 kDa	

Applications

Tested Applications: FC, WB, ELISA	Positive Controls: WB : Hela cells, human placenta tissue
Species Specificity: human	

Background Information

GM2A, also named as GM2-AP and SAP-3, is the large binding pocket which can accommodate several single chain phospholipids and fatty acids. GM2A also exhibits some calcium-independent phospholipase activity. It plays a key role in the degradation of ganglioside GM2 (GM2). GM2A stimulates only the breakdown of ganglioside GM2 and glycolipid GA2 by beta-hexosaminidase A.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of GM2A antibody (66080-1-Ig; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GM2A transfected HeLa cells.



HeLa cells were subjected to SDS PAGE followed by western blot with 66080-1-Ig (GM2A antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



1X10⁶ HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human GM2A (66080-1-Ig, Clone:1E4D3) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).