

For Research Use Only

NSE Monoclonal antibody

Catalog Number: 66150-1-Ig

Featured Product

16 Publications



Basic Information

Catalog Number: 66150-1-Ig	GenBank Accession Number: BC002745	Purification Method: Protein A purification
Size: 150ul , Concentration: 2093 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2026	CloneNo.: 6F8G3
Source: Mouse	Full Name: enolase 2 (gamma, neuronal)	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:2500-1:10000 IF 1:50-1:400
Isotype: IgG1	Calculated MW: 47 kDa	
Immunogen Catalog Number: AG19106	Observed MW: 47 kDa	

Applications

Tested Applications:
FC, IF, IHC, WB, ELISA

Cited Applications:
IF, IHC, WB

Species Specificity:
human, mouse, rat, pig

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : PC-12 cells, C6 cells, fetal human brain tissue, HEK-293 cells, Neuro-2a cells, pig brain tissue, rat brain tissue, SH-SY5Y cells, U-251 cells, pig cerebellum tissue, rat cerebellum tissue, mouse brain tissue, mouse cerebellum tissue

IHC : mouse brain tissue, human brain tissue, rat cerebellum tissue, rat brain tissue

IF : SH-SY5Y cells, mouse brain tissue

Background Information

NSE, also named as ENO2, belongs to the enolase family. Enolases are cytoplasmic glycolytic enzymes that may be involved in differentiation. The enolase has three isoenzymes, alpha, beta and gamma. The alpha form is expressed in most tissues, whereas the beta form is expressed in muscle tissue. The gamma enolase (ENO2), a homodimer, is primarily localized in neurons and neuroendocrine cells and is a cancer diagnostic marker for brain tumors (PMID:7520111). ENO2 plays a role in the glycolysis-related energy pathway and might be involved in higher metabolic activity during the day than at night, at least in part.

Notable Publications

Author	Pubmed ID	Journal	Application
Rongkun Li	34836938	Cell Death Dis	WB
Huan Liu	36377337	Brain Behav	IF
Jiaren Pan	36419844	Dis Markers	WB,IHC

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66150-1-Ig (NSE antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using NSE antibody (66150-1-Ig, Clone: 6F8G3) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




1×10^6 SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human NSE (66150-1-Ig, Clone: 6F8G3) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).




WB result of NSE antibody (66150-1-Ig; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NSE transfected U-251 cells.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 66150-1-Ig (NSE antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 66150-1-Ig (NSE antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of SH-SY5Y cells using 66150-1-Ig (NSE antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG (H+L).