For Research Use Only

CoraLite®594-conjugated HDAC1 Monoclonal antibody

Catalog Number: CL594-66085



Purification Method:

Excitation/Emission maxima

wavelengths:

588 nm / 604 nm

Basic Information

Catalog Number: GenBank Accession Number:

CL594-66085 BC000301 Protein G purification
Size: GeneID (NCBI): CloneNo.:

100ul , Concentration: 1000 ug/ml by 3065 4C4G7
Nanodrop:

Nanodrop; UNIPROT ID: Recommended Dilutions: Source: Q13547 IF/ICC: 1:4000-1:16000

Mouse Full Name: FC (Intra): 0.20 ug per 10^6 cells in a

Isotype: histone deacetylase 1
IgG1 Calculated MW:

IgG1 Calculated MW:
Immunogen Catalog Number: 55 kDa

AG19128 Observed MW:

60-65 kDa

Applications Tested Applications: Positive Controls:

IF/ICC, FC (Intra)

Species Specificity:
human, mouse, rat

IF/ICC : A431 cells, HeLa cells
FC (Intra) : HeLa cells,

Background Information

Histone deacetylases(HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues leading to the formation of a condensed and transcriptionally silenced chromatin. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family and is a component of the histone deacetylase complex, which is responsible for gene expression silencing. It also plays an important role in the control of cell proliferation and differentiation by interacting with RB, p53 and other transcription factors. At least 4 classes of HDAC were identified. As a class I HDAC, HDAC 1 was primarily found in the nucleus. This antibody is raised against residues near the C terminus of human HDAC1. The calculated molecular weight of HDAC1 is 55 kDa, but modified HDAC1 is about 60-65 kDa.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% glycerol, 0.05% Proclin300, 0.5% BSA, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CL594-66085 (HDAC1 antibody) at dilution of 1:100.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug CoraLite®594 Anti-Human HDAC1 (CL594-66085, Clone:4C4G7) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using Coralite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite®594 HDAC1 antibody (CL594-66085, Clone: 4C4G7) at dilution of 1:8000.