For Research Use Only

G3BP1 Polyclonal antibody

Catalog Number: 13057-2-AP

Featured Product

157 Publications



Basic Information

Catalog Number: GenBank Accession Number:

13057-2-AP BC006997 GeneID (NCBI): Size: 150ul , Concentration: 650 $\mu g/ml$ by 10146

Nanodrop: **UNIPROT ID:** Q13283 Rabbit Full Name:

Isotype: GTPase activating protein (SH3 domain) binding protein 1 IgG

Immunogen Catalog Number: Calculated MW:

466 aa, 52 kDa AG3728 Observed MW: 68 kDa

Tested Applications:

WB, IP, IF, FC, IHC, ELISA Cited Applications:

WB, IP, IF, RIP, IHC, CoIP Species Specificity: human, rat, mouse

Cited Species:

human, chicken, rat, mouse, monkey, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC 1:200-1:800 IF 1:1000-1:4000

Applications

Positive Controls:

WB: C6 cells, HEK-293 cells, human brain tissue. Neuro-2a cells, HeLa cells, HepG2 cells, MCF-7 cells, Jurkat cells, mouse kidney tissue, rat kidney tissue, mouse brain tissue, rat brain tissue

IP: HEK-293 cells,

IHC: human colon cancer tissue, human breast cancer tissue, human lung cancer tissue

IF: sodium arsenite treated HeLa cells.

Background Information

GAP SH3 Binding Protein 1 (G3BP1), also named as G3BP, is an effector of stress granule (SG) assembly. SG biology plays an important role in the pathophysiology of TDP-43 in ALS and FTLD-U. G3BP1 can be used as a marker of SG. It has been shown to function downstream of Ras and play a role in RNA metabolism, signal transduction, and proliferation. G3BP1 is a ubiquitously expressed protein that localizes to the cytoplasm in proliferating cells and to the nucleus in non-proliferating cells. G3BP1 has recently been implicated in cancer biology.

Notable Publications

Author	Pubmed ID	Journal	Application
Jumin Park	34551427	Nucleic Acids Res	IF
Bin Dai	34560101	J Biol Chem	IF
Jozsef Gal	31481451	Mol Cell Biol	WB,IF

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free

in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13057-2-AP (G3BP1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours




Immunofluorescent analysis of (4% PFA) fixed sodium arsenite treated HeLa cells using G3BP1 antibody (13057-2-AP) at dilution of 1:2000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




WB result of G3BP1 antibody (13057-2-AP; 1:100000; incubated at room temperature for 1.5 hours) with sh-Control and sh-G3BP1 transfected HFK-93 rells




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 13057-2-AP (G3BP1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 13057-2-AP (G3BP1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-G3BP1 (IP:13057-2-AP, 4ug; Detection:13057-2-AP 1:1000) with HEK-293 cells lysate 1040 ug.



1X10^6 HeLa cells were intracellularly stained with 0.2 ug Anti-Human G3BP1 (13057-2-AP) and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).