For Research Use Only

TNNC1 Polyclonal antibody

Catalog Number: 13504-1-AP

Featured Product

6 Publications



Basic Information

Catalog Number: GenBank Accession Number: 13504-1-AP BC030244

Size: GeneID (NCBI):

150ul , Concentration: 1000 μ g/ml by 7134 Nanodrop and 573 μ g/ml by Bradford Full Name:

method using BSA as the standard; troponin C type 1 (slow)

Source: Calculated MW:
Rabbit 161 aa, 18 kDa

Isotype: Observed MW:
IgG 18 kDa

Immunogen Catalog Number:

AG4339

Purification Method: Antigen affinity purification Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:500-1:1000

for WB IHC 1:50-1:500

IF

Applications

Tested Applications:

IF, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human, mouse, rat Cited Species:

human, mouse, rat

Positive Controls:

WB: mouse skeletal muscle tissue, human stomach

tissue

IP: mouse heart tissue,

IHC: human skeletal muscle tissue, mouse skeletal

muscle tissue

IF: U2OS,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

Troponin is a central regulatory protein of striated muscle contraction, and together with tropomyosin, is located on the actin filament. Troponin consists of 3 subunits: Tnl, which is the inhibitor of actomyosin ATPase; TnT, which contains the binding site for tropomyosin; and TnC, the protein encoded by this gene. TnC plays an important role in excitation-contraction coupling and is the primary target of compounds such as levosimendan that augment Ca2+ responsiveness of the thin filament.

Notable Publications

Author	Pubmed ID	Journal	Application
Magdalena A Cichewicz	30037979	Mol Cell Biol	WB
Xian Ye	32946432	Med Sci Monit	WB
Fan Ma	33116603	Onco Targets Ther	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data




mouse skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 13504-1-AP (TNNC1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human skeletal muscle tissue slide using 13504-1-AP (TNNC1 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human skeletal muscle tissue slide using 13504-1-AP (TNNC1 Antibody) at dilution of 1:200 (under 40x lens).



IP Result of anti-TNNC1 (IP:13504-1-AP, 4ug; Detection:13504-1-AP 1:600) with mouse heart tissue lysate 4400ug.



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using 13504-1-AP (TNNC 1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).