

For Research Use Only

TAF7 Polyclonal antibody

Catalog Number: 13506-1-AP



Basic Information

Catalog Number: 13506-1-AP	GenBank Accession Number: BC032737	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 700 µg/ml by Nanodrop and 367 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 6879	Recommended Dilutions: WB 1:1000-1:6000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
Source: Rabbit	Full Name: TAF7 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 55kDa	IHC 1:20-1:200
Isotype: IgG	Calculated MW: 349 aa, 40 kDa	
Immunogen Catalog Number: AG4406	Observed MW: 55 kDa	

Applications

Tested Applications: IHC, IP, WB, ELISA	Positive Controls:
Species Specificity: human, mouse, rat	WB : Jurkat cells, HeLa cells, MCF-7 cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IP : Jurkat cells,
	IHC : human cervical cancer tissue,

Background Information

TAF7, also named as Transcription initiation factor TFIID 55 kDa subunit, is a 349 amino acid protein, which belongs to the TAF7 family. TAF7 functions as a component of the DNA-binding general transcription factor complex TFIID, a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. TAF7 is phosphorylated by CIITA. The calculated molecular weight of TAF7 is 40 kDa, but phosphorylated TAF7 is about 55kDa. (PMID: 22711989)

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 13506-1-AP (TAF7 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 13506-1-AP (TAF7 antibody) at dilution of 1:100 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human cervical cancer using 13506-1-AP (TAF7 antibody) at dilution of 1:100 (under 40x lens).



IP Result of anti-TAF7 (IP:13506-1-AP, 4ug; Detection:13506-1-AP 1:500) with Jurkat cells lysate 1200ug.