For Research Use Only

ENY2 Polyclonal antibody

Catalog Number: 15778-1-AP

Featured Product

1 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method: Antigen affinity purification

15778-1-AP Size:

BC007870 GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 340 µg/ml by

56943

WB 1:500-1:1000

Nanodrop and 340 µg/ml by Bradford Full Name:

method using BSA as the standard;

enhancer of yellow 2 homolog (Drosophila)

IHC 1:50-1:500

Rabbit

Calculated MW:

Isotype: IgG

12 kDa Observed MW:

Immunogen Catalog Number:

10 kDa

AG8462

Applications

Tested Applications:

Positive Controls:

IHC, WB, ELISA

WB: PC-3 cells

Cited Applications:

IHC: human breast cancer tissue,

IP, WB

Species Specificity:

human

Cited Species:

buffer pH 6.0

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

Background Information

ENY2, also named as DC6, is involved in mRNA export coupled transcription activation by association with both the TREX-2 and the SAGA complexes. Within the SAGA complex, participates in a subcomplex that specifically deubiquitinates both histones H2A and H2B. The SAGA complex is recruited to specific gene promoters by activators such as MYC, where it is required for transcription. It is required for nuclear receptor-mediated transactivation. The TREX-2 complex functions in docking export-competent ribonucleoprotein particles (mRNPs) to the nuclear entrance of the nuclear pore complex (nuclear basket). The MW of ENY2 is about 10 kDa.

Notable Publications

Author Pubmed ID Journal Application S Vijayalingam 26559831 J Virol WB. IP

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 15778-1-AP (ENY2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



PC-3 cells were subjected to SDS PAGE followed by western blot with 15778-1-AP (ENY2 Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.