

For Research Use Only

# NMD3 Polyclonal antibody

Catalog Number: 16060-1-AP **5 Publications**



## Basic Information

<b>Catalog Number:</b> 16060-1-AP	<b>GenBank Accession Number:</b> BC013317	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 260 µg/ml by Nanodrop and 267 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 51068	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB
<b>Source:</b> Rabbit	<b>Full Name:</b> NMD3 homolog (S. cerevisiae)	<b>IHC 1:50-1:500</b>
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 503 aa, 58 kDa	<b>IF 1:50-1:500</b>
<b>Immunogen Catalog Number:</b> AG8894	<b>Observed MW:</b> 58 kDa	

## Applications

<b>Tested Applications:</b> IF, IHC, IP, WB, ELISA	<b>Positive Controls:</b>
<b>Cited Applications:</b> IF, WB	<b>WB :</b> HepG2 cells, L02 cells, mouse liver tissue, rat liver tissue
<b>Species Specificity:</b> human, mouse, rat	<b>IP :</b> HepG2 cells,
<b>Cited Species:</b> human, mouse	<b>IHC :</b> human breast cancer tissue, human liver tissue
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	<b>IF :</b> HeLa cells,

## Background Information

Human NMD3 gene encodes 60S ribosomal export protein which was found in a 60S ribosomal subunit export complex with RAN and XPO1. This nucleocytoplasmic shuttling protein NMD3 is an adaptor for export of the 60S ribosomal subunit from the nucleus. NMD3 contains a CRM-1-dependent leucine-rich nuclear export signal (NES) and a dispersed nuclear localization signal (NLS), the basic region of which is also required for nucleolar accumulation. NMD3 are required for nuclear export of the 60S ribosomal subunit in yeast and vertebrate cells, recent finding has also revealed its role in Arabidopsis thaliana.

## Notable Publications

Author	Pubmed ID	Journal	Application
Wong Chi C CC	21803848	Blood	WB
Andrew J Finch	21536732	Genes Dev	WB
Kaosheng Lv	33711283	Cell Stem Cell	WB

## Storage


**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 16060-1-AP (NMD3 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




IP Result of anti-NMD3 (IP:16060-1-AP, 3ug; Detection:16060-1-AP 1:800) with HepG2 cells lysate 1720ug.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 16060-1-AP (NMD3 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 16060-1-AP (NMD3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 16060-1-AP (NMD3 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).