

For Research Use Only

PIP4K2C Polyclonal antibody

Catalog Number: 17077-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number: 17077-1-AP	GenBank Accession Number: BC028596	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 450 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 79837	Recommended Dilutions: WB 1:1000-1:4000 IHC 1:50-1:500
Source: Rabbit	Full Name: phosphatidylinositol-5-phosphate 4-kinase, type II, gamma	
Isotype: IgG	Calculated MW: 421 aa, 47 kDa	
Immunogen Catalog Number: AG10989	Observed MW: 47 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls: WB : HeLa cells, HepG2 cells, PC-12 cells
Cited Applications: WB	IHC : human liver cancer tissue,
Species Specificity: human, mouse, rat	
Cited Species: human	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

PIP4K2C is involved in several processes, including 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate biosynthetic process; negative regulation of 1-phosphatidylinositol-4-phosphate 5-kinase activity; and positive regulation of autophagosome assembly. PIP4Ks negatively regulate insulin signaling through a catalytic-independent mechanism.

Notable Publications

Author	Pubmed ID	Journal	Application
Diana G Wang	31091439	Cell Rep	WB
Ismael AL-Ramahi	29256861	Elife	WB
Ron Benyair	37064812	Autophagy Rep	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 17077-1-AP (PIP4K2C antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 17077-1-AP (PIP4K2C antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 17077-1-AP (PIP4K2C antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 17077-1-AP (PIP4K2C antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 17077-1-AP (PIP4K2C antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 17077-1-AP (PIP4K2C antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.