For Research Use Only

## IL-5 Polyclonal antibody

Catalog Number:17786-1-AP



Basic Information	Catalog Number: 17786-1-AP	GenBank Accession Number: BC066282	Purification Method: Antigen affinity purification
	Size: 150ul, Concentration: 260 µg/ml by Bradford method using BSA as the standard; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG11840	GenelD (NCBI): 3567	WB 1:1000-1:4000 olony-stimulating nil)
		UNIPROT ID: P05113 Full Name: interleukin 5 (colony-stimulating	
		Calculated MW: 134 aa, 15 kDa	
		Applications	
Species Specificity: human			
Background Information	Interleukin-5, or IL-5, was originally discovered as a soluble T cell-derived factor, called T cell-replacing factor (TRF), that induced T cell-depleted activated B cells to secrete immunoglobulin. IL-5 is a key hematopoietic cytokine in eosinophil differentiation, maturation, recruitment and activation at sites of allergic inflammation. IL-5 also plays a role in the development, metabolism, and function of basophils. IL-5 exerts its biological activity through the IL-5 receptor (IL-5R), which is composed of at least two chains: an α chain that binds IL-5 with low affinity and a β chain that does not bind IL-5, but together with the IL-5α chain, constitutes the high affinity IL-5 receptor. The β chain is common to the IL-3, IL-5 and GM-CSF receptors and has been shown to signal through the JAK/Stat pathway. IL-5 has long been associated with the cause of several allergic diseases including allergic rhinitis and asthma, wherein a large increase in the number of circulating, airway tissue, and induced sputum eosinophils have been observed.		
	and a β chain that does not bind IL-5, β chain is common to the IL-3, IL-5 an pathway. IL-5 has long been associat asthma, wherein a large increase in t	ch is composed of at least two chains: but together with the IL-5a chain, con d GM-CSF receptors and has been sho ed with the cause of several allergic of	stitutes the high affinity IL-5 receptor. The wn to signal through the JAK/Stat diseases including allergic rhinitis and
Storage	and a β chain that does not bind IL-5, β chain is common to the IL-3, IL-5 an pathway. IL-5 has long been associat asthma, wherein a large increase in t	ch is composed of at least two chains: but together with the IL-5a chain, con d GM-CSF receptors and has been sho ed with the cause of several allergic of the number of circulating, airway tissue the shipment.	stitutes the high affinity IL-5 receptor. The wn to signal through the JAK/Stat diseases including allergic rhinitis and

For technical support and original validation data for this product please contact:T: 1 (888) 4PTGLAB (1-888-478-4522) (toll freeE: proteintech@ptglab.comin USA), or 1(312) 455-8498 (outside USA)W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Recombinant protein were subjected to SDS PAGE followed by western blot with 17786-1-AP (IL5 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.