# For Research Use Only

# IL-3 Polyclonal antibody

Catalog Number: 17835-1-AP

1 Publications



**Basic Information** 

Catalog Number: 17835-1-AP

GenBank Accession Number: BC066273

Antigen affinity purification

Size:

GeneID (NCBI):

Recommended Dilutions:

150ul, Concentration: 500 µg/ml by Nanodrop and 347 µg/ml by Bradford

3562

WB 1:1000-1:4000

**Purification Method:** 

IHC 1:50-1:500

method using BSA as the standard;

interleukin 3 (colony-stimulating

factor, multiple)

Rabbit Isotype: Calculated MW:

152 aa, 17 kDa IgG Observed MW:

20 kDa

Immunogen Catalog Number: AG11839

**Applications** 

**Tested Applications:** 

IHC. WR.FIISA

**Cited Applications:** 

WB: Recombinant protein, Recombinant protein

protein

Positive Controls:

IHC: human tonsillitis tissue,

Species Specificity: human

WB

**Cited Species:** 

pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

# **Background Information**

Interleukin-3 (IL-3) is a multilineage hematopoietic growth factor that promotes the proliferation, differentiation and survival of early multilineage hematopoietic progenitors. In particular, this cytokine plays a key role in stimulating the proliferation and survival of myeloid precursors. It is involved in a variety of cell activities such as cell growth, differentiation and apoptosis. This cytokine has been shown to also possess neurotrophic activity, and it may be associated with neurologic disorders (PMID: 24103757;2698876;2809196)

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Dwight D Harris	37808040	JTCVS Open	WB

### Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


## **Selected Validation Data**



Recombinant protein were subjected to SDS PAGE followed by western blot with 17835-1-AP (IL3 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 17835-1-AP (IL3 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 17835-1-AP (IL3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).