

For Research Use Only

AXUD1 Polyclonal antibody

Catalog Number: 18162-1-AP

Featured Product

2 Publications



Basic Information

Catalog Number:

18162-1-AP

Size:

150ul , Concentration: 450 µg/ml by Nanodrop and 260 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG6158

GenBank Accession Number:

BC038949

GeneID (NCBI):

64651

Full Name:

AXIN1 up-regulated 1

Calculated MW:

64 kDa

Observed MW:

64-70 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

IF, IHC, WB

Species Specificity:

human, mouse, rat

Cited Species:

mouse, rat

Positive Controls:

WB : HeLa cells, HEK-293 cells, mouse thymus tissue

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Xing Ye	28830684	Exp Cell Res	WB, IF, IHC
Yan Li	37077375	Heliyon	WB, IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 18162-1-AP (AXUD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 18162-1-AP (AXUD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



mouse thymus tissue were subjected to SDS PAGE followed by western blot with 18162-1-AP (AXUD1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.