

For Research Use Only

NR5A1 Polyclonal antibody

Catalog Number: 18658-1-AP **13 Publications**



Basic Information

Catalog Number: 18658-1-AP	GenBank Accession Number: BC032501	Purification Method: Antigen affinity purification
Size: 150ul, Concentration: 500 µg/ml by Nanodrop and 300 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 2516	Recommended Dilutions: WB 1:500-1:1000 IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC 1:50-1:500
Source: Rabbit	Full Name: nuclear receptor subfamily 5, group A, member 1	
Isotype: IgG	Calculated MW: 52 kDa	
Immunogen Catalog Number: AG13252	Observed MW: 52 kDa	

Applications

Tested Applications:
FC, IHC, IP, WB, ELISA

Cited Applications:
IF, IHC, WB

Species Specificity:
human, mouse, rat

Cited Species:
camel, human, goat, mouse, Ondatra zibethicus

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: mouse ovary tissue, rat ovary tissue

IP: A2780 cells,

IHC: human ovary tissue, human liver tissue, rat testis tissue

Background Information

Steroidogenic factor-1 (SF-1, STF-1), also known as NR5A1, regulates multiple genes involved in the adrenal and gonadal development and in the biosynthesis of a variety of hormones, including adrenal and gonadal steroids, anti-Mullerian hormone (AMH), and gonadotropins. SF-1 belongs to the fushi tarazu factor-1 (FTZ-F1) subfamily of orphan nuclear receptors. Initially identified as a tissue-specific transcriptional regulator of cytochrome P450 steroid hydroxylases, research studies of both global and tissue-specific knockout mice have demonstrated that SF-1 is required for the development of adrenal glands, gonads, ventromedial hypothalamus, and for the proper functioning of pituitary gonadotropes. Indeed, humans with mutations that render SF-1 transcriptionally inactive can present with testicular failure, ovarian failure, and adrenal insufficiency. Furthermore, dysregulation of SF-1 has been linked to diseases such as endometriosis and adrenocortical carcinoma. Like other nuclear hormone receptors, SF-1 has a modular domain structure composed of an N-terminal zinc finger DNA-binding domain, a ligand-binding domain, a C-terminal AF-2 activation domain, and a hinge region with AF-1-like activation activity. SF-1 also contains a fushi tarazu factor 1 box, which functions as an accessory DNA binding domain. SF-1 is primarily phosphorylated at Ser203, which is thought to enhance its transcriptional activity by promoting complex formation with transcriptional cofactors. In addition to phosphorylation at Ser203, SF-1 is subject to SUMO conjugation and acetylation at ε-amino groups of target lysine residues. Whereas SUMOylation represses SF-1 function, acetylation enhances its transcriptional activity. In the adult ovary, SF-1 localizes to theca/interstitial cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Wenqian Xie	32991988	J Steroid Biochem Mol Biol	IHC
Haibo Zhang	34671938	Reprod Sci	IHC
Jianlin Liang	31710289	Elife	IF

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

***** 20ul sizes contain 0.1% BSA**


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 18658-1-AP (NR5A1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



1×10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human NR5A1 (18658-1-AP) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



IP Result of anti-NR5A1 (IP:18658-1-AP, 4ug; Detection:18658-1-AP 1:300) with A2780 cells lysate 960ug.



Immunohistochemical analysis of paraffin-embedded human ovary tissue slide using 18658-1-AP (NR5A1 Antibody) at dilution of 1:100 (under 40x lens).