

CCDC27 Polyclonal antibody

Catalog Number: 23769-1-AP

Basic Information

Catalog Number: 23769-1-AP	GenBank Accession Number: BC112108	Purification Method: Antigen Affinity purified
Size: 150ul , Concentration: 500 µg/ml by Nanodrop and 340 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 148870	Recommended Dilutions: WB 1:500-1:1000 IHC 1:50-1:500
Source: Rabbit	Full Name: coiled-coil domain containing 27	
Isotype: IgG	Calculated MW: 656 aa, 75 kDa	
Immunogen Catalog Number: AG20676	Observed MW: 75 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Species Specificity: human	WB : human testis tissue, IHC : human kidney tissue,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information


CCDC27 also named as coiled coil domain containing 27 is a 656 amino acids protein, which contains a coiled-coil domain. CCDC27 expression is predominantly in male testis and lowly in other tissues. CCDC27 is a candidate gene for low bone mineral density. The function of CCDC27 is still non-known and need to be further studied.

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



human testis tissue were subjected to SDS PAGE followed by western blot with 23769-1-AP (CCDC27 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 23769-1-AP (CCDC27 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).