For Research Use Only

BDNF Polyclonal antibody

Catalog Number: 25699-1-AP

Featured Product

29 Publications



Basic Information

Catalog Number: GenBank Accession Number:

25699-1-AP BC029795
Size: Genel D (NCBI):

150ul , Concentration: 300 µg/ml by 627

Nanodrop; UNIPROT ID:
Source: P23560
Rabbit Full Name:

Isotype: brain-derived neurotrophic factor

IgGCalculated MW:Immunogen Catalog Number:247 aa, 28 kDaAG22489Observed MW:

30 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications: IF. IHC. WB

Species Specificity: human, rat, mouse Cited Species:

human, rat, mouse, macaque

Positive Controls:

WB: mouse brain tissue, mouse cerebellum tissue, rat

Purification Method:

WB 1:500-1:1000

Antigen affinity purification

Recommended Dilutions:

brain tissue

Background Information

BDNF is a member of the family of neurotrophic factors. It was first explored and purified from pig brain in 1982, and other members of the neurotrophin family were successively discovered. BDNF is induced by cortical neurons, and is necessary for survival of striatal neurons in the brain. Expression of BDNF is reduced in both Alzheimer's and Huntington disease patients. It participates in axonal growth, pathfinding and in the modulation of dendritic growth and morphology. It also plays a role in the regulation of stress response and in the biology of mood disorders. The BDNF gene is rather complex in its structure and it can express five different mRNA isoforms by alternative splicing, producing 28-37 kDa proteins.

Notable Publications

Author	Pubmed ID	Journal	Application
John H Brock	26414795	J Neurotrauma	IHC
Wu WenBo	28952006	Neurochem Res	WB,IHC
Chenxia Sheng	29057264	Biomed Res Int	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 25699-1-AP (BDNF antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.