For Research Use Only

BMP6 Polyclonal antibody

Catalog Number: 55421-1-AP



Basic Information

Catalog Number: GenBank Accession Number:

55421-1-AP NM_001718
Size: GeneID (NCBI):

150ul, Concentration: 800 µg/ml by 654

anodrop;

Source: bone morphogenetic protein 6

Rabbit Calculated MW:
Isotype: 57 kDa
IgG Observed MW:
57 kDa

Purification Method: Antigen affinity purification Recommended Dilutions: WB 1:500-1:1000

Applications

Tested Applications:

WB, ELISA

Species Specificity: human, mouse

Positive Controls:

WB: A549 cells, mouse liver tissue, HEK-293 cells, HL-

60 cells, U2OS cells

Background Information

Bone morphogenetic protein 6 (BMP6) is also named as VG-1-related protein (VG-1-R, VGR-1) and VGR, and belongs to the TGF-beta family. (PMID:33878365). Growth factor of the TGF-beta superfamily plays essential roles in many developmental processes including cartilage and bone formation (PMID:31019025). BMP6 participates in iron homeostasis, inhibits invasion by increasing adhesions and cell-cell type interactions and induces angiogenesis directly on vascular endothelial cells. BMP6 is coded by a tumor suppressor gene whose subexpression is related to the development and cancer progression; during neoplastic processes, methylation is the main mechanism by which gene silencing occurs (PMID:34706618). BMP6 plays an important role in the regulation of hepcidin/HAMP expression and iron metabolism by acting as a ligand for hemojuvelin/HJV (PMID:26582087). BMP6 initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2B (PMID:18070108).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.


Storage Buffe

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 55421-1-AP (BMP6 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.



mouse liver tissue were subjected to SDS PAGE followed by western blot with 55421-1-AP (BMP6 Antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.