

## Beta Actin Monoclonal antibody

Catalog Number: 60008-1-Ig

2945 Publications

## Basic Information

## Catalog Number:

60008-1-Ig

## Size:

150ul, Concentration: 1000 µg/ml by 60  
Bradford method using BSA as the  
standard;

## Source:

Mouse

## Isotype:

IgM

## Immunogen Catalog Number:

AG0297

## GenBank Accession Number:

BC002409

## GeneID (NCBI):

60

## UNIPROT ID:

P60709

## Full Name:

actin, beta

## Calculated MW:

375 aa, 42 kDa

## Observed MW:

42 kDa

## Purification Method:

Caprylic acid/ammonium sulfate  
precipitation

## CloneNo.:

7D2C10

## Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:50-1:500

## Applications

## Tested Applications:

WB, FC, IHC, ELISA

## Cited Applications:

WB, IP, IF, IHC, CoIP

## Species Specificity:

human, mouse, rat, plant, Zebrafish

## Cited Species:

cat, human, goat, tick, chicken, yeast, rat, Arabidopsis,  
Terminalia bellirica, Rare minnow

**Note-IHC: suggested antigen retrieval with  
TE buffer pH 9.0; (\*) Alternatively, antigen  
retrieval may be performed with citrate  
buffer pH 6.0**

## Positive Controls:

**WB** : multi-cells/tissue, HeLa cells, MCF-7 cells, HEK-  
293 cells, A549 cells, rice whole plant tissue,  
arabidopsis whole plant tissue

**IHC** : human colon tissue, human brain tissue, human  
kidney tissue, human lung tissue, human spleen  
tissue, human ovary tissue, human testis tissue,  
human placenta tissue, human heart tissue

## Background Information

Beta actin, also named as ACTB and F-Actin, belongs to the actin family. Actins are highly conserved globular proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells. At least six isoforms of actins are known in mammals and other vertebrates: alpha (ACTC1, cardiac muscle 1), alpha 1 (ACTA1, skeletal muscle) and 2 (ACTA2, aortic smooth muscle), beta (ACTB), gamma 1 (ACTG1) and 2 (ACTG2, enteric smooth muscle). Beta and gamma 1 are two non-muscle actin proteins. Most actins consist of 376aa, while ACTG2 (rich in muscles) has 375aa and ACTG1(found in non-muscle cells) has only 374aa. Beta actin has been widely used as the internal control in RT-PCR and Western Blotting as a 42-kDa protein. However, the 37-40 kDa cleaved fragment of beta actin can be generated during apoptosis process. This antibody can recognize all the actins.

## Notable Publications

Author	Pubmed ID	Journal	Application
Qiaoxia Zheng	36198318	Cell	WB
Menghui Xu	36290677	Antioxidants (Basel)	WB
BreAnna Cameron	34590699	Biol Open	WB

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech  
Group brand and is not available to purchase from any  
other manufacturer.**


Selected Validation Data




Western blot analysis of Beta-actin in various tissues and cell lines using Proteintech antibody 60008-1-Ig at a dilution of 1:5000. Extra bands were detected in some species with unknown reason.




Immunohistochemical analysis of paraffin-embedded human colon using 60008-1-Ig(ACTB antibody) at dilution of 1:50 (under 10x lens).




Immunohistochemical analysis of paraffin-embedded human colon using 60008-1-Ig(ACTB antibody) at dilution of 1:50 (under 40x lens).



Western blot of HeLa cell with anti-Actin-Beta (60008-1-Ig) at various dilutions.



Arabidopsis whole plant tissue were subjected to SDS PAGE followed by western blot with 60008-1-Ig (beta actin Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



1X10<sup>6</sup> HepG2 cells were stained with 0.2ug beta actin antibody (60008-1-Ig, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L) with dilution 1:1000.