For Research Use Only

AHCY Monoclonal antibody

Catalog Number:66019-1-lg 1 Publications



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

66019-1-lg

BC011606 GeneID (NCBI): Protein A purification

Size: 150ul , Concentration: 2900 μ g/ml by 191 CloneNo.:

Nanodrop and 1300 µg/ml by Bradford_{UNIPROT ID:}

1F9B10

method using BSA as the standard;

P23526

Recommended Dilutions: WB 1:500-1:2000

Source: Mouse

Full Name:

IHC 1:20-1:200 IF 1:2000-1:8000

WB: HeLa cells, human testis tissue, LO2 cells, MCF-7

IHC: human testis tissue, human placenta tissue

Isotype:

S-adenosylhomocysteine hydrolase Calculated MW:

48 kDa

lgG2b Immunogen Catalog Number:

Observed MW:

AG18036

48 kDa

Applications

Tested Applications:

WB, IF, IHC, ELISA

Cited Applications:

WB

Positive Controls:

Species Specificity:

human

IF: HeLa cells,

Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

AHCY, also named as SAHH, belongs to the adenosylhomocysteinase family. AHCY is a competitive inhibitor of Sadenosyl-L-methionine-dependent methyl transferase reactions; therefore AHCY may play a key role in the control of methylations via regulation of the intracellular concentration of adenosylhomocysteine. AHCY catalyzes the reversible hydrolysis of S-adenosylhomocysteine to adenosine and L-homocysteine. (PMID:21220475).

Notable Publications

Author **Pubmed ID** Journal Application Yankun Wang Cell Death Differ WB 35974250

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


Selected Validation Data

 $150 \text{kd} \rightarrow$ $100 \text{kd} \rightarrow$ $70 \text{kd} \rightarrow$ $50 \text{kd} \rightarrow$ $40 \text{kd} \rightarrow$ $30 \text{kd} \rightarrow$


HeLa cells were subjected to SDS PAGE followed by western blot with 66019-1-1g (AHCY antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human testis using 66019-1-lg(AHCY antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human testis using 66019-1-Ig(AHCY antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using AHCY antibody (66019-1-lg, Clone: 1F9B10) at dilution of 1:4000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).