

Catalog Number: HZ-1019

Data Sheet





Animal Component-Free

Human cell expressed

Tag-Free


Endotoxin Free

Product Description


Animal-free Recombinant Human IL-6 is produced in human 293 cells as a monomeric glycoprotein with an apparent molecular mass of 21 to 29 kDa. This broad molecular mass is attributable to glycosylation, which is absent when this cytokine is expressed in E. coli. Glycosylation contributes to stability in cell growth media and other applications. IL-6 is a pleiotropic cytokine that plays an important role in host defense by regulating immune and inflammatory responses. Produced by T cells, monocytes, fibroblasts, endothelial cells, and keratinocytes, IL-6 has diverse biological functions.

Alternative Names	B cell stimulatory factor 2, BSF 2, BSF2, CDF, CTL differentiation factor, HGF, HSF, Hybridoma growth factor, IFN beta 2, IFNB2, IL 6, IL-6, Interferon beta 2, Interleukin 6
Source	Human Embryonic Kidney cells (HEK293). HEK293-derived IL-6 protein
Species Reactivity	human,mouse,monkey,rabbit,rat

Specifications					
Test	Method	Specification			
Activity	Dose-dependent stimulation of the proliferation of 3G12B10 cells (mouse hybridoma cell line)	0.03-0.24 ng/mL EC50 in 3G12B10 cells.			
Molecular Mass	SDS-PAGE	24 and 29 kDa reduced, 21 and 28 kDa non-reduced, monomer, glycosylated			
Purity	SDS-PAGE	>95%			
Endotoxin	LAL	<1 EU/µg			



Recombinant human IL-6 (HZ-1019) stimulates dosedependent proliferation of the 3G12B10 hybridoma cell line. Cell number was quantitatively assessed by PrestoBlue® Cell Viability Reagent. 3G12B10 cells were treated with increasing concentrations of recombinant IL-6 for 96 hours. The EC50 range is 0.03-0.24 ng/mL.



www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

Preparation				
Shipping Temperature				
Formulation	Formulation 1x PBS, See Certificate of Analysis for details			
Reconstitution Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile 1xPBS pH 7.4 corendotoxin-free recombinant human serum albumin (HSA).				

	Product Form	Temperature Conditions	Storage Time (From Date of Receipt)
	Lyophilized	-20°C to -80°C	Until Expiry Date
Stability and Storage	Lyophilized	Room Temperature	2 weeks
_	Reconstituted as per CofA	-20°C to -80°C	6 months
	Reconstituted as per CofA	4°C	1 week
		Avoid repeated freeze-thaw cycles.	

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc.

5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com