

For Research Use Only

NOB1 Polyclonal antibody

Catalog Number: 10091-2-AP

Featured Product

7 Publications



Basic Information

Catalog Number:

10091-2-AP

Size:

150UL, Concentration: 450 µg/ml by Nanodrop and 180 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0141

GenBank Accession Number:

BC000050

GeneID (NCBI):

28987

Full Name:

NIN1/RPN12 binding protein 1 homolog (S. cerevisiae)

Calculated MW:

47 kDa

Observed MW:

47-50 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:200-1:1000

for WB

IHC 1:10-1:100

Applications

Tested Applications:

IHC, IP, WB, ELISA

Cited Applications:

IHC, WB

Species Specificity:

human

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, SGC-7901 cells, HepG2 cells, K-562 cells

IP: HepG2 cells,

IHC: human lymphoma tissue, human placenta tissue, human colon cancer tissue

Background Information

NOB1 was first identified in yeast as an essential gene encoding the Nin one binding protein, which is involved in pre-rRNA processing. In a late cytoplasmic processing step, Nob1 cleaves a 20S rRNA intermediate at cleavage site D to produce the mature 18S rRNA. In addition, NOB1 is a crucial molecule in the maturation of the 20S proteasome and protein degradation. It serves as a chaperone to join the 20S proteasome with the 19S regulatory particle in the nucleus and facilitates the maturation of the 20S proteasome. Recently NOB1 has been reported to be overexpressed in several types of cancer, suggesting its involvement in the tumorigenesis.

Notable Publications

Author	Pubmed ID	Journal	Application
Sabina Schütz	30201955	Nat Commun	WB
Xin Gao	27035645	Oncol Rep	
Xiao-Wen He	25624720	World J Gastroenterol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 10091-2-AP (NOB1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lymphoma using 10091-2-AP (NOB1 antibody) at dilution of 1:25 (under 10x lens).



IP Result of anti-NOB1 (IP:10091-2-AP, 4ug; Detection:10091-2-AP 1:300) with HepG2 cells lysate 2400ug.