### For Research Use Only

# PRDX3 Polyclonal antibody

Catalog Number: 10664-1-AP

Featured Product

34 Publications



**Basic Information** 

Catalog Number: GenBank Accession Number:

10664-1-AP BC007062

Size: GeneID (NCBI):

150ul, Concentration: 600 ug/ml by 10935

Nanodrop; UNIPROT ID:

Source: P30048
Rabbit Full Name:
Isotype: peroxiredoxin 3
IgG Calculated MW:

Immunogen Catalog Number: 27 kDa

AG1062 Observed MW:

26-28 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions:

WB: 1:5000-1:50000 IHC: 1:50-1:500 IF/ICC: 1:200-1:800

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, ELISA

**Cited Applications:** 

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, MCF-7 cells, LNCaP cells, mouse brain tissue, mouse heart tissue, rat brain

tissue

IHC: human lung cancer tissue,

IF/ICC: HepG2 cells,

## **Background Information**

PRDX3(Peroxiredoxin 3), also named as AOP1, HBC189 and MER5, belongs to the ahpC/TSA family. It is involved in redox regulation of the cell and protects radical-sensitive enzymes from oxidative damage by a radical-generating system. PRDX3 is required for MYC-mediated proliferation, transformation, and apoptosis after glucose withdrawal and essential for maintaining mitochondrial mass and membrane potential in transformed rat and human cells (PMID:12011429). PRDX3 acts synergistically with MAP3K13 to regulate the activation of NF-kappa-B in the cytosol.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Ling Fu	31411056	Antioxid Redox Signal	WB
Caitlin E. Miller	36358489	Antioxidants (Basel)	WB
Eirini Meimaridou	29046340	J Endocrinol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3  $\,$ 

Aliquoting is unnecessary for -20°C storage


\*\*\* 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 10664-1-AP (PRDX3 antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 10664-1-AP (PRDX3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 10664-1-AP (PRDX3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



HeLa cells were subjected to SDS PAGE followed by western blot with 10664-1-AP (PRDX3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using PRDX3 antibody (10664-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



WB result of PRDX3 antibody (10664-1-AP; 1:12000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PRDX3 transfected HEK-293 cells.