

OPTN Polyclonal ANTIBODY

Catalog Number: 10837-1-AP

Featured Product

33 Publications

Basic Information

Catalog Number:
10837-1-AP

Size:
52 µg/150 µl

Source:
Rabbit

Isotype:
IgG

Purification Method:
Antigen affinity purification

Immunogen Catalog Number:
AG1272

GenBank Accession Number:
BC013876

GeneID (NCBI):
10133

Full Name:
optineurin

Calculated MW:
66 kDa

Observed MW:
66; 70-74 kDa

Recommended Dilutions:

WB 1:2000-1:12000

IP 0.5-4.0 µg for IP and 1:500-1:1000 for WB

IHC 1:50-1:500

IF 1:10-1:100

Applications

Tested Applications:

FC, IF, IHC, IP, WB, ELISA

Cited Applications:

IF, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : HeLa cells; mouse testis tissue, mouse brain tissue

IP : mouse brain tissue;

IHC : human gliomas tissue; human pancreas cancer tissue

IF : MCF-7 cells;

Background Information

OPTN, also named as FIP2, GLC1E, HIP7, HYPL and NRP, plays a neuroprotective role in the eye and optic nerve. It is probably part of the TNF-alpha signaling pathway that can shift the equilibrium toward induction of cell death. OPTN may act by regulating membrane trafficking and cellular morphogenesis via a complex that contains Rab8 and huntingtin (HD). OPTN may constitute a cellular target for adenovirus E3 14.7, an inhibitor of TNF-alpha functions, thereby affecting cell death. Defects in OPTN are the cause of primary open angle glaucoma type 1E (GLC1E). Defects in OPTN are a cause of susceptibility to normal pressure glaucoma (NPG). OPTN mutated in adult-onset primary open angle glaucoma. It supports the protective role of OPTN in the trabecular meshwork. OPTN has 3 isoforms with MW 66, 65 and 60 kDa (refer to UniProt). Catalog#10837-1-AP recognises 66 and 70-74 kDa band, and the additional 70-74 kDa band due to phosphorylation.

Notable Publications

Author	Pubmed ID	Journal	Application
Madhu Puri	28872615	Toxins (Basel)	WB
Hongyu Li	30160596	Autophagy	WB
Norihiko Furuya	30309841	EMBO Rep	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




various lysates were subjected to SDS PAGE followed by western blot with 10837-1-AP (OPTN antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours




IP Result of anti-OPTN (IP:10837-1-AP, 3ug; Detection:10837-1-AP 1:800) with mouse brain tissue lysate 8000ug.




Immunofluorescent analysis of MCF-7 cells, using OPTN antibody 10837-1-AP at 1:25 dilution and Rhodamine-labeled goat anti-rabbit IgG (red).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 10837-1-AP (OPTN Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 10837-1-AP (OPTN Antibody) at dilution of 1:200 (under 40x lens).



1X10⁶ MCF-7 cells were stained with 0.2ug OPTN antibody (10837-1-AP, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1500.