

For Research Use Only

# SCNN1A Polyclonal antibody

Catalog Number: 10924-2-AP

Featured Product

14 Publications



## Basic Information

<b>Catalog Number:</b> 10924-2-AP	<b>GenBank Accession Number:</b> BC006526	<b>Purification Method:</b> Antigen affinity purification
<b>Size:</b> 150ul , Concentration: 187 µg/ml by Bradford method using BSA as the standard;	<b>GeneID (NCBI):</b> 6337	<b>Recommended Dilutions:</b> WB 1:500-1:1000 IHC 1:20-1:200 IF 1:50-1:500
<b>Source:</b> Rabbit	<b>Full Name:</b> sodium channel, nonvoltage-gated 1 alpha	
<b>Isotype:</b> IgG	<b>Calculated MW:</b> 76 kDa	
<b>Immunogen Catalog Number:</b> AG1325	<b>Observed MW:</b> 60-70 kDa	

## Applications

<b>Tested Applications:</b> IF, IHC, WB, ELISA	<b>Positive Controls:</b> WB : PC-13 cells, HEK-293 IHC : human kidney tissue, IF : A549 cells,
<b>Cited Applications:</b> IHC, WB	
<b>Species Specificity:</b> human, mouse, rat	
<b>Cited Species:</b> human, mouse, rat	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

SCNN1A (sodium channel, non-voltage-gated 1 alpha), also known as ENaC (epithelial Na<sup>+</sup>) channel subunit alpha) or amiloride-sensitive sodium channel subunit alpha, is the alpha subunit of the epithelial Na<sup>+</sup> channel (ENaC). ENaC is expressed in the apical membrane of salt-absorbing epithelia of kidney, distal colon, and lung. ENaC is a non-voltage gated, constitutively active channel highly selective for sodium. It has an essential role in salt and fluid homeostasis across epithelial tissues. ENaC consists of three different subunits: alpha, beta, gamma. Mutations in the gene of SCNN1A have been associated with pseudohypoaldosteronism type 1 (PHA1), a rare salt wasting disease resulting from target organ unresponsiveness to mineralocorticoids. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. It has been reported that full-length SCNN1A protein can be cleaved into 65- and 30-kDa fragments (PMID: 16477034; 18701608).

## Notable Publications

Author	Pubmed ID	Journal	Application
Ivana d'Angelo	29035132	J Aerosol Med Pulm Drug Deliv	WB
Jianping Zhang	31719660	Sci Rep	IHC
David W Scott	28481660	Am J Respir Crit Care Med	WB


## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)  
E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




WB result of SCNN1A (10924-2-AP, 1:1000) with si-control and si-SCNN1A transfected HEK293 cells.




PC-13 cells were subjected to SDS PAGE followed by western blot with 10924-2-AP (SCNN1A Antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 10924-2-AP (SCNN1A Antibody) at dilution of 1:50 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 10924-2-AP (SCNN1A Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed A549 cells using 10924-2-AP (SCNN1A antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).