

For Research Use Only

# RANGAP1 Polyclonal antibody

Catalog Number: 11102-1-AP

1 Publications



## Basic Information

**Catalog Number:**

11102-1-AP

**Size:**

150ul , Concentration: 500 µg/ml by Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG1563

**GenBank Accession Number:**

BC014044

**GeneID (NCBI):**

5905

**Full Name:**

Ran GTPase activating protein 1

**Calculated MW:**

64 kDa

**Observed MW:**

64 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:500-1:2000

IP 0.5-4.0 ug for IP and 1:200-1:1000 for WB

IHC 1:50-1:500

IF 1:50-1:500

## Applications

**Tested Applications:**

FC, IF, IHC, IP, WB, ELISA

**Cited Applications:**

WB

**Species Specificity:**

human

**Cited Species:**

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

WB : SH-SY5Y cells, HEK-293 cells

IP : HEK-293 cells,

IHC : human gliomas tissue,

IF : HepG2 cells,

## Background Information

### Notable Publications

Author	Pubmed ID	Journal	Application
Xianwei Chen	26844761	J Proteomics	WB

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 11102-1-AP (RANGAP1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 11102-1-AP (RANGAP1 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 11102-1-AP (RANGAP1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 11102-1-AP (RANGAP1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10<sup>6</sup> HepG2 cells were stained with 0.2ug RANGAP1 antibody (11102-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.



IP Result of anti-RANGAP1 (IP:11102-1-AP, 4ug; Detection:11102-1-AP 1:300) with HEK-293 cells lysate 3600ug.