

For Research Use Only

MYH9 Polyclonal antibody

Catalog Number: 11128-1-AP

Featured Product

50 Publications



Basic Information

Catalog Number:

11128-1-AP

Size:

150ul, Concentration: 900 ug/ml by Nanodrop;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1612

GenBank Accession Number:

BC011915

GeneID (NCBI):

4627

UNIPROT ID:

P35579

Full Name:

myosin, heavy chain 9, non-muscle

Calculated MW:

227 kDa

Observed MW:

224 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB: 1:5000-1:50000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC: 1:50-1:500

IF-P: 1:50-1:500

IF/ICC: 1:50-1:500

FC (Intra): 0.25 ug per 10⁶ cells in a 100 µl suspension

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, IP, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, RIP

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, chicken

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A549 cells, mouse brain tissue, mouse kidney tissue, mouse colon tissue, human brain tissue, U2OS cells

IP: mouse brain tissue,

IHC: human lung cancer tissue, human kidney tissue, mouse kidney tissue

IF-P: mouse small intestine tissue,

IF/ICC: HepG2 cells,

FC (Intra): HepG2 cells,

Background Information

MYH9 (Myosin-9) appears to play a role in cytokinesis, cell shape, and specialized functions such as secretion and capping. MYH9 is a hexameric protein that consists of two heavy chains (~200 kDa) and two pairs of non-identical light chains (15-22 kDa), a pair of essential or alkali light chains, and a pair of regulatory light chains (PMID: 12800156).

Notable Publications

Author	Pubmed ID	Journal	Application
Peiyuan Li	34586803	J Agric Food Chem	WB
Yang Cao	30336116	Exp Cell Res	WB,IP,IF
Ru Zheng	27742864	Reproduction	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various cell lysates were subjected to SDS PAGE followed by western blot with 11128-1-AP (MYH9 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 11128-1-AP (MYH9 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 11128-1-AP (MYH9 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-MYH9 (IP:11128-1-AP, 5ug; Detection:11128-1-AP 1:1000) with mouse brain tissue lysate 4000ug.




Immunohistochemical analysis of paraffin-embedded human normal colon slide using 11128-1-AP (MYH9 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse small intestine tissue using MYH9 antibody (11128-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using MYH9 antibody (11128-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red).



1×10^6 HepG2 cells were intracellularly stained with 0.25 ug MYH9 Polyclonal antibody (11128-1-AP) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).