

For Research Use Only

# ERK1/2 Polyclonal antibody

Catalog Number: 11257-1-AP

Featured Product

422 Publications



## Basic Information

### Catalog Number:

11257-1-AP

### Size:

150ul, Concentration: 500 ug/ml by Nanodrop;

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG1759

### GenBank Accession Number:

BC013992

### GeneID (NCBI):

5595

### UNIPROT ID:

P27361

### Full Name:

mitogen-activated protein kinase 3

### Calculated MW:

43 kDa

### Observed MW:

38-44 kDa

### Purification Method:

Antigen affinity purification

### Recommended Dilutions:

WB 1:2000-1:16000

IHC 1:100-1:400

IF/ICC 1:200-1:800

## Applications

### Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

### Cited Applications:

WB, IHC, IF, ColP

### Species Specificity:

human, mouse, rat

### Cited Species:

human, mouse, rat, pig, canine, chicken, zebrafish, sheep

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

### Positive Controls:

**WB** : NIH/3T3 cells, HeLa cells, HEK-293 cells, PC-12 cells, C6 cells, mouse brain tissue, mouse lung tissue

**IHC** : human colon cancer tissue, human breast cancer tissue

**IF/ICC** : HEK-293 cells,

## Background Information

ERK1 and ERK2 belongs to the protein kinase superfamily. It is involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK-1. ERK1/2 catalyzed the reaction: ATP + a protein = ADP + a phosphoprotein. It is activated by tyrosine phosphorylation in response to INS and NGF. This antibody can recognize both ERK1 and ERK2 with the molecular mass of 38-44 kDa.

## Notable Publications

Author	Pubmed ID	Journal	Application
Xiao-Feng Zhu	36180975	Phytother Res	WB
Hui Wang	36243659	Hepatobiliary Pancreat Dis Int	WB
Hongxia Li	34558311	Epigenomics	WB

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

\*\*\* 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11257-1-AP (ERK1/2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of ERK1/2 antibody (11257-1-AP, 1:1000) with HeLa and HEK293 cells.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11257-1-AP (ERK1/2 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11257-1-AP (ERK1/2 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug Anti-Human ERK1/2 (11257-1-AP) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using ERK1/2 antibody (11257-1-AP) at dilution of 1:400 and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).