For Research Use Only

Rab18 Polyclonal antibody

Catalog Number:11304-1-AP

Featured Product 17 Publications




Basic Information	Catalog Number: 11304-1-AP	GenBank Accession Number: BC015014		Purification Method: Antigen affinity purification	
	Size: 150ul , Concentration: 450 ug/ml by Nanodrop; Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG1841	GeneID (NCBI): y 22931 UNIPROT ID: Q9NP72 Full Name: RAB18, member RAS Calculated MW: 206 aa, 23 kDa Observed MW: 20-23 kDa	oncogene family	Recommended Dilutions: WB: 1:1000-1:6000 IP: 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate IHC: 1:200-1:2000 IF-P: 1:50-1:500 IF/ICC: 1:200-1:800	
Applications	Tested Applications:		Positive Controls:		
	WB, IHC, IF/ICC, IF-P, IP, ELISA Cited Applications: WB, IHC, IF			tissue, human heart tissue, human cells, human testis tissue, mouse rain tissue	
	human, mouse, rat IHC : hun		IP: mouse test	tis tissue, rat testis tissue	
			IHC : human stomach cancer tissue, mouse brain tissue		
	Cited Species: human, mouse, pig, monkey		IF-P : mouse brain tissue,		
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IF/ICC : HEK-2	93 cells,	
Background Information	and may be implicated in transport	between the plasma m	embrane and ear	nrole in apical endocytosis/recycling, ly endosomes. It also plays a key role ir ssociated with Warburg micro syndrome	
Notable Publications	Author Pr	ubmed ID Jour	nal	Application	
			t Cell Dev Biol	IHC	
			Targets Ther	WB,IHC	
	5		Targets Ther	WB,IHC	
Storage	Storage: Store at -20°C. Stable for one year a Storage Buffer: PBS with 0.02% sodium azide and ! Aliquoting is unnecessary for -20°C	50% glycerol, pH7.3			


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 11304-1-AP (Rab18 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




human testis tissue were subjected to SDS PAGE followed by western blot with 11304-1-AP (RAB18 antibody) at dilution of 1:800 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 11304-1-AP (Rab18 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using Rab18 antibody (11304-1-AP) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Rab18 antibody (11304-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using Rab18 antibody (11304-1-AP) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IP result of anti-Rab18 (IP:11304-1-AP, 4ug; Detection:11304-1-AP 1:2000) with mouse testis tissue lysate 1120 ug.



IP result of anti-Rab18 (IP:11304-1-AP, 4ug; Detection:11304-1-AP 1:2000) with rat testis tissue lysate 4000 ug.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 11304-1-AP (Rab18 antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).