

For Research Use Only

BubR1 Polyclonal antibody

Catalog Number: 11504-2-AP

Featured Product

8 Publications



Basic Information

Catalog Number: 11504-2-AP	GenBank Accession Number: BC018739	Purification Method: Antigen affinity purification
Size: 150UL, Concentration: 247 µg/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 701	Recommended Dilutions: WB 1:500-1:2000 IHC 1:100-1:400 IF 1:20-1:200
Source: Rabbit	Full Name: budding uninhibited by benzimidazoles 1 homolog beta (yeast)	
Isotype: IgG	Calculated MW: 1050 aa, 120 kDa	
Immunogen Catalog Number: AG2053	Observed MW: 120-130 kDa	

Applications

Tested Applications: IF, IHC, WB, ELISA	Positive Controls: WB: mouse testis tissue, HeLa cells, human testis tissue, K-562 cells IHC: human colon cancer tissue, IF: SKOV-3 cells,
Cited Applications: IF, IHC, WB	
Species Specificity: human, mouse	
Cited Species: human, mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

BUB1B (Mitotic checkpoint serine/threonine-protein kinase BUB1 beta), also named as BUBR1, MAD3 and SSK1, belongs to the BUB1 subfamily. Human BUB1B is a 120 kDa protein that plays a central role in the spindle assembly checkpoint (SAC), the evolutionary conserved and self-regulatory system of higher organisms that monitors and repairs defects in chromosome segregation in mitotic cells (PMID:22201036). In addition to phosphorylation, BUB1B is also subjected to posttranslational modifications including acetylation (PMID:22167194). Defects in BUB1B are the cause of premature chromatid separation trait (PCS) and mosaic variegated aneuploidy syndrome type 1 (MVA1).

Notable Publications

Author	Pubmed ID	Journal	Application
Zhi-Chuan Zhu	29520102	Oncogene	WB
Fengjie Yuan	31291454	Nucleic Acids Res	WB
Wei Xu	32210727	Int J Biol Sci	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse testis tissue were subjected to SDS PAGE followed by western blot with 11504-2-AP (BubR1 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.




WB result of BubR1 antibody (11504-2-AP; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-BubR1 transfected HeLa cells.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11504-2-AP (BubR1 antibody at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 11504-2-AP (BubR1 antibody at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of SKOV-3 cells using 11504-2-AP (BubR1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).