

For Research Use Only

GOSR2/Membrin Polyclonal antibody



Catalog Number: 12095-1-AP

Featured Product

Basic Information

Catalog Number:

12095-1-AP

GenBank Accession Number:

BC009710

Purification Method:

Antigen affinity purification

Size:

150UL, Concentration: 1000 µg/ml by 9570

GeneID (NCBI):

9570

Recommended Dilutions:

WB 1:500-1:3000

Nanodrop and 333 µg/ml by Bradford method using BSA as the standard;

Full Name:

golgi SNAP receptor complex member

IF 1:50-1:500

Source:

Rabbit

2

Calculated MW:

212 aa, 25 kDa

Isotype:

IgG

Observed MW:

25 kDa

Immunogen Catalog Number:

AG2736

Applications

Tested Applications:

IF, WB, ELISA

Positive Controls:

WB : Jurkat cells, human liver tissue, mouse liver tissue, mouse stomach tissue, LO2 cells, rat liver tissue

Species Specificity:

human, mouse, rat

IF : HepG2 cells,

Background Information

Golgi SNAP receptor complex member 2 (GOSR2), also known as GS27 or Membrin, is a member of the soluble NSF attachment protein receptor (SNARE) family of vesicle docking proteins. GOSR2 is localized in Golgi apparatus. It is involved in transport of proteins from the cis/medial-Golgi to the trans-Golgi network.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of GOSR2/Membrin antibody (12095-1-AP; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GOSR2/Membrin transfected Jurkat cells.



Various lysates were subjected to SDS PAGE followed by western blot with 12095-1-AP (GOSR2/Membrin antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 12095-1-AP (GOSR2/Membrin antibody) at dilution of 1:50 and CoraLite488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).