

For Research Use Only

NARF Polyclonal antibody

Catalog Number:12173-1-AP



Basic Information

Catalog Number: 12173-1-AP	GenBank Accession Number: BC016440	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 133 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 26502	Recommended Dilutions: WB 1:500-1:3000
Source: Rabbit	UNIPROT ID: Q9UHQ1	
Isotype: IgG	Full Name: nuclear prelamin A recognition factor	
Immunogen Catalog Number: AG2816	Calculated MW: 46 kDa, 53 kDa	
	Observed MW: 52 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human liver tissue, human lung tissue, human placenta tissue, mouse testis tissue
Species Specificity: human, mouse, rat	

Background Information

Prelamin A, is processed in the nucleus to lamin A by removal of its final 18 amino acids, including the cysteine residue in its C-terminal CAAX box, which is farnesylated. Nuclear prelamin A recognition factor (NARF) binds the farnesylated prelamin A C-terminal domain(preAct)(PMID:10514485). NARF, at 52 kDa, also named as IOP2, is a small enough protein to freely diffuse through the nuclear pore complex and may be retained within the nucleus by binding nuclear proteins and it has all of the information necessary to be imported into the nucleus(PMID:10514485). It has 3 isoforms produced by alternative splicing with the molecular mass of 51 kDa, 56 kDa and 46 kDa, respectively.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human liver tissue were subjected to SDS PAGE followed by western blot with 12173-1-AP (NARF antibody) at dilution of 1:400 incubated at room temperature for 1.5 hours.