

For Research Use Only

CREB1 Polyclonal ANTIBODY



Catalog Number: 12208-1-AP

Featured Product

19 Publications

Basic Information

Catalog Number:

12208-1-AP

Size:

150UL, Concentration: 287 µg/ml by Bradford method using BSA as the standard;

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG2852

GenBank Accession Number:

BC010636

GeneID (NCBI):

1385

Full Name:

cAMP responsive element binding protein 1

Calculated MW:

341 aa, 35 kDa

Observed MW:

43-46 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:500-1:3000

IP 0.5-4.0 ug for IP and 1:500-1:1000 for WB

IHC 1:50-1:500

IF 1:10-1:100

Applications

Tested Applications:

FC, IF, IHC, IP, WB, ELISA

Cited Applications:

ChIP, IHC, WB

Species Specificity:

human, mouse, rat, monkey

Cited Species:

human, mouse, rat, sea cucumbers

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: multi-cells/tissue, COS-7 cells, HEK-293 cells, HeLa cells, Jurkat cells, K-562 cells, mouse brain tissue, mouse lung tissue, NIH/3T3 cells, rat brain tissue

IP: HEK-293 cells,

IHC: human cervical cancer tissue, mouse brain tissue, human prostate cancer tissue, human thyroid tissue

IF: HEK-293 cells,

Background Information

CREB1, also named as CREB, belongs to the bZIP family, containing one bZIP domain and one KID (kinase-inducible) domain. This protein binds the cAMP response element (CRE), a sequence present in many viral and cellular promoters. CREB stimulates transcription on binding to the CRE. This protein is stimulated by phosphorylation. Phosphorylation of both Ser-133 and Ser-142 in the SCN regulates the activity of CREB and participates in circadian rhythm generation. Phosphorylation of Ser-133 allows CREBBP binding. Transcription activation is enhanced by the TORC coactivators which act independently of Ser-133 phosphorylation. CREB1 is sumoylated by SUMO1. Sumoylation on Lys-304, but not on Lys-285, is required for nuclear localization of this protein. Sumoylation is enhanced under hypoxia, promoting nuclear localization and stabilization. Defects in CREB1 may be a cause of angiomatoid fibrous histiocytoma (AFH), a distinct variant of malignant fibrous histiocytoma that typically occurs in children and adolescents and is manifest by nodular subcutaneous growth. A chromosomal aberration involving CREB1 is found in a patient with angiomatoid fibrous histiocytoma. Translocation t(2;22)(q33;q12) with CREB1 generates a EWSR1/CREB1 fusion gene that is most common genetic abnormality in this tumor type. CREB1 exists some isoforms and range of calculated molecular weight of isoforms are 35-37 kDa and 25 kDa, but the modified CREB1 protein is about 43 kDa (PMID: 25883219).

Notable Publications

Author	Pubmed ID	Journal	Application
Chenxia Sheng	29057264	Biomed Res Int	WB
Disi Bai	30542609	Toxicol Res (Camb)	WB
Xiqiong Han	30459620	Front Pharmacol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 12208-1-AP (CREB1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




WB result of 12208-1-AP (CREB1 antibody) with various lysates at dilution of 1:1,500.



IP Result of anti-CREB1 (IP:12208-1-AP, 3ug; Detection:12208-1-AP 1:600) with HEK-293 cells lysate 2000ug.



Immunofluorescent analysis of HEK-293 cells using 12208-1-AP (CREB1 antibody) at dilution of 1:25 and Rhodamine-Goat anti-Rabbit IgG.



1×10^6 HEK-293 cells were stained with 0.2ug CREB1 antibody (12208-1-AP, red) and control antibody (blue). Fixed with 90% MeOH blocked with 3% BSA (30 min). Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) with dilution 1:1000.