

For Research Use Only

# ZFP36L1/2 Polyclonal ANTIBODY



Catalog Number: 12306-1-AP

Featured Product

1 Publications

## Basic Information

**Catalog Number:**

12306-1-AP

**Size:**

150UL, Concentration: 247 µg/ml by Bradford method using BSA as the standard;

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG2952

**GenBank Accession Number:**

BC018340

**GeneID (NCBI):**

677

**Full Name:**

zinc finger protein 36, C3H type-like 1

**Calculated MW:**

338 aa, 36 kDa

**Observed MW:**

45-50 kDa

**Purification Method:**

Antigen affinity purification

**Recommended Dilutions:**

WB 1:500-1:1000

IHC 1:20-1:200

IF 1:20-1:200

## Applications

**Tested Applications:**

IF, IHC, WB, ELISA

**Cited Applications:**

IF

**Species Specificity:**

human, mouse, rat

**Cited Species:**

human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

WB: U-937 cells, 3T3-L1 cells, RAW 264.7 cells, HeLa cells

IHC: human placenta tissue, human ovary tissue

IF: HepG2 cells,

## Background Information

ZFP36L1, also named as Butyrate response factor 1 or TPA-induced sequence 11b, is a 338 amino acid protein, which is phosphorylated by RPS6KA1 at Ser-334 upon phorbol 12-myristate 13-acetate (PMA) treatment. ZFP36 encodes tristetraprolin (TTP) the prototype of a small family of RBPs, called the ZFP36 family, that are characterised by highly conserved tandem CCCH zinc-finger RNA-binding domains [PMID: 10751406]. ZFP36 is a RBP that promotes RNA decay and negatively regulates the expression of the myogenic regulatory factor MyoD by binding to the 3'UTR of MyoD mRNA [PMID: 25815583]. Mouse satellite cells from Zfp36-deficient mice express increased amounts of MyoD and display impaired satellite activation, demonstrating a role for ZFP36 in the maintenance of quiescence [PMID: 25815583]. The functions of the ZFP36L1 and ZFP36L2 family members have not been evaluated in skeletal muscle stem cell fate, but have been shown to act redundantly to promote quiescence during lymphocyte development. ZFP36L1 has been implicated in the persistence of the marginal zone B lymphocyte population. ZFP36L1 exists three isoform through blasting on NCBI database and can be phosphorylated, so the range of the molecular weight of ZFP36L1 is about 40-50 kDa. (PMID: 26180518, PMID: 17030620). This antibody recognizes both ZFP36L1 and ZFP36L2.

## Notable Publications

Author	Pubmed ID	Journal	Application
Weirui Ma	30449617	Cell	IF

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




U-937 cells were subjected to SDS PAGE followed by western blot with 12306-1-AP (ZFP36L1 antibody) at dilution of 1:600 incubated at room temperature for 1.5 hours.




WB result of ZFP36L1/2 antibody (12306-1-AP; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-ZFP36L1/2 transfected HeLa cells.



Immunohistochemical analysis of paraffin-embedded human ovary tissue slide using 12306-1-AP (ZFP36L1 Antibody) at dilution of 1:50 (under 40x lens).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 12306-1-AP (ZFP36L1 Antibody) at dilution of 1:50 (under 40x lens).



Immunofluorescent analysis of HepG2 cells using 12306-1-AP (ZFP36L1 antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).