For Research Use Only

CA8 Polyclonal antibody

Catalog Number: 12391-1-AP

Featured Product

9 Publications



Basic Information

Catalog Number: GenBank Accession Number: 12391-1-AP BC015531

12391-1-AP BC015531
Size: GeneID (NCBI):

150ul , Concentration: 650 ug/ml by 767

Nanodrop and 333 ug/ml by Bradford UNIPROT ID: method using BSA as the standard; P35219

Source: Full Name:

Rabbit carbonic anhydrase VIII

Isotype: Calculated MW:
IgG 290 aa, 32 kDa
Immunogen Catalog Number: Observed MW:
AG3068 32 kDa

Purification Method: Antigen affinity purification

Recommended Dilutions: WB: 1:2000-1:12000

IP: 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IHC: 1:50-1:500

Applications

Tested Applications:

WB, IP, IHC, ELISA

Cited Applications: WB, IHC, IF, IP

Species Specificity:

human, mouse, rat Cited Species:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, mouse brain tissue, Jurkat cells, K-

562 cells

IP: mouse brain tissue,

IHC: human gliomas tissue, mouse cerebellum tissue

Background Information

The CA8 (CARP) gene encodes carbonic anhydrase VIII, which is part of a family of zinc metalloenzyme. CA8 has a central carbonic anhydrase motif, but it lacks carbonic anhydrase activity due to absence of catalytic zinc coordinating residues(PMID:2121526). CARP is a novel IP3R1-binding protein, and is expressed in Purkinje cells abundantly. CA8 is co-localized with IP3R1 in Purkinje cells and it binds to IP3R1, reducing the affinity of the receptor for its ligand, IP3(PMID:12611586). Defects in CA8 are the cause of cerebellar ataxia mental retardation and dysequilibrium syndrome type 3 (CMARQ3)(PMID:19461874).

Notable Publications

Author	Pubmed ID	Journal	Application
Huai-Lu Ma	31715371	Biomed Pharmacother	WB,IHC
Ken Asada	33928345	Nucleic Acids Res	IP
Liddelow Shane A SA	23843944	PLoS One	IHC

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


*** 20ul sizes contain 0.1% BSA

For technical support and original validation data for this product please contact:


T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com W: ptglab.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 12391-1-AP (CA8 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 12391-1-AP (CA8 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-CA8 (IP:12391-1-AP, 4ug; Detection:12391-1-AP 1:1000) with mouse brain tissue lysate 4000ug.