

For Research Use Only

ARNT2 Polyclonal antibody

Catalog Number:12810-1-AP

Featured Product

3 Publications



Basic Information

Catalog Number: 12810-1-AP	GenBank Accession Number: BC036099	Purification Method: Antigen affinity purification
Size: 150ul , Concentration: 260 ug/ml by Nanodrop and 227 ug/ml by Bradford method using BSA as the standard;	GeneID (NCBI): 9915	Recommended Dilutions: WB: 1:500-1:1000 IHC: 1:50-1:500
Source: Rabbit	UNIPROT ID: Q9HBZ2	
Isotype: IgG	Full Name: aryl-hydrocarbon receptor nuclear translocator 2	
Immunogen Catalog Number: AG3536	Calculated MW: 716 aa, 79 kDa	
	Observed MW: 80-85 kDa	

Applications

Tested Applications: WB, IHC, ELISA	Positive Controls: WB : HEK-293 cells, Jurkat cells, Raji cells IHC : human gliomas tissue,
Cited Applications: WB	
Species Specificity: human, mouse, rat	
Cited Species: human, mouse	

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Aryl-hydrocarbon receptor nuclear translocator 2 (ARNT2), is a member of the basic-helix-loop-helix-Per-Arnt-Sim (bHLH-PAS) superfamily of transcription factors. It specifically recognizes the xenobiotic response element (XRE). ARNT2 plays an important role in normal glucose handling in pancreatic beta cell function in humans and mice. ARNT2 proteins are dimeric partners for other PAS proteins such as HIF and SIM. ARNT2 and the other ARNT family members, e.g., ARNT, hypoxin inducible factor 1 α (HIF1 α), HIF2 α , and aryl hydrocarbon receptor (AhR), are thought to function as heterodimers, and regulate the expression, and thereby the function, of many genes. Under hypoxic conditions, it complexes with hypoxia-inducible factor 1 α in the nucleus and this complex binds to hypoxia-responsive elements in enhancers and promoters of oxygen-responsive genes. Modulation of ARNT2 in pancreatic beta cells affects glucose stimulated INS secretion (GSIS). This antibody is a rabbit polyclonal antibody raised against residues near the C terminus of human ARNT2.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhixiao Sun	33537006	Front Microbiol	WB
Wen-Juan Li	40173041	Cell Rep	WB
Maralice Conacci-Sorrell	39184078	Res Sq	WB

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

*** 20ul sizes contain 0.1% BSA


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



WB result of ARNT2 antibody (12810-1-AP; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ARNT2 transfected Jurkat cells.



Immunohistochemical analysis of paraffin-embedded human gliomas tissue slide using 12810-1-AP (ARNT2 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0)).



HEK-293 cells were subjected to SDS PAGE followed by western blot with 12810-1-AP (ARNT2 antibody) at dilution of 1:300 incubated at room temperature for 1.5 hours.